

Effect of Lowering the Dialysate Temperature in Chronic Hemodialysis: A Systematic Review and Meta-Analysis

Reem A. Mustafa,^{*†} Fadi Bdair,[†] Elie A. Akl,^{*‡} Amit X. Garg,^{*§||} Heather Thiesen-Philbrook,[¶] Hassan Salameh,[†] Sood Kisra,[†] Jihad Nesrallah,^{**} Ahmad Al-Jaishi,[§] Parth Patel,^{††} Payal Patel,^{††} Ahmad A. Mustafa,^{§§} and Holger J. Schünemann^{*|||}

Abstract

Background and objectives Lowering the dialysate temperature may improve outcomes for patients undergoing chronic hemodialysis. We reviewed the reported benefits and harms of lower temperature dialysis.

Design, setting, participants, & measurements We searched the Cochrane Central Register, OVID MEDLINE, EMBASE, and Pubmed until April 15, 2015. We reviewed the reference lists of relevant reviews, registered trials, and relevant conference proceedings. We included all randomized, controlled trials that evaluated the effect of reduced temperature dialysis versus standard temperature dialysis in adult patients receiving chronic hemodialysis. We followed the Grading of Recommendations Assessment, Development and Evaluation approach to assess confidence in the estimates of effect (*i.e.*, the quality of evidence). We conducted meta-analyses using random effects models.

Results Twenty-six trials were included, consisting of a total of 484 patients. Compared with standard temperature dialysis, reduced temperature dialysis significantly reduced the rate of intradialytic hypotension by 70% (95% confidence interval, 49% to 89%) and significantly increased intradialytic mean arterial pressure by 12 mmHg (95% confidence interval, 8 to 16 mmHg). Symptoms of discomfort occurred 2.95 (95% confidence interval, 0.88 to 9.82) times more often with reduced temperature compared with standard temperature dialysis. The effect on dialysis adequacy was not significantly different, with a Kt/V mean difference of -0.05 (95% confidence interval, -0.09 to 0.01). Small sample sizes, loss to follow-up, and a lack of appropriate blinding in some trials reduced confidence in the estimates of effect. None of the trials reported long-term outcomes.

Conclusions In patients receiving chronic hemodialysis, reduced temperature dialysis may reduce the rate of intradialytic hypotension and increase intradialytic mean arterial pressure. High-quality, large, multicenter, randomized trials are needed to determine whether reduced temperature dialysis affects patient mortality and major adverse cardiovascular events.

Clin J Am Soc Nephrol 11: 442–457, 2016. doi: 10.2215/CJN.04580415

Introduction

Approximately 2 million patients worldwide receive life-sustaining chronic hemodialysis treatments (1). Intradialytic hypotension occurs in $\leq 30\%$ of hemodialysis treatments and is associated with substantial long-term mortality (2–4). Maggiore *et al.* (5) proposed cool dialysis to prevent intradialytic hypotension by increasing peripheral vascular resistance, improving cardiac output, and altering the levels of vasoactive peptides. Nephrologists, however, remain cautious about using this modality of hemodialysis because of limited evidence about its long-term effects and the assumed disadvantages. These disadvantages include the discomfort that patients may experience in association with cool dialysis and the concern that it may interfere with achieving adequate dialysis.

In 2006, Selby and McIntyre (6) conducted a systematic review to assess the effects of lowering

dialysate temperature. The review was restricted to English language articles, and Selby and McIntyre (6) were unable to report the effects of lowering dialysate temperature on long-term patient-important outcomes because of lack of data. Since 2006, more trials have been published on this topic (7,8). We conducted this systematic review to evaluate the effect of cool dialysis compared with standard temperature dialysis on patient-important outcomes in adults undergoing maintenance chronic hemodialysis.

Materials and Methods

Data Sources and Searches

We conducted this systematic review in accordance with a prespecified registered protocol available at <http://www.crd.york.ac.uk/PROSPERO> (registration no. CRD42011001104). We reported the results

Departments of

*Clinical Epidemiology and Biostatistics and

|||Medicine, McMaster University, Hamilton, Ontario, Canada;

†Departments of Medicine and Biomedical and Health Informatics and

††School of Medicine, University of Missouri, Kansas City, Missouri;

‡Department of Internal Medicine, American University of Beirut, Beirut, Lebanon;

§Department of Medicine, London Health Sciences Centre, London, Ontario, Canada;

||Department of Medicine, Western University, London, Ontario, Canada;

¶Program of Applied Translational Research, Department of Internal Medicine, Yale

University School of Medicine, New Haven, Connecticut; **Li Ka Shing Knowledge

Institute, St. Michael's Hospital, Toronto, Ontario, Canada; ††MD

Program, St. James School of Medicine, Chicago, Illinois; and

§§School of Medicine, Jordan University of Science and Technology, Erbil, Jordan

Correspondence:

Dr. Reem A. Mustafa, Department of Clinical Epidemiology and Biostatistics, McMaster University, HSC Room 2C15, 1280 Main Street, West Hamilton, ON, Canada L8S 4K1. Email: ramustafa@gmail.com

according to Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines (9).

We performed an electronic search of the Cochrane Central Register of Controlled Trials (until April of 2015), OVID MEDLINE (from 1946 to April of 2015), EMBASE (from 1947 to April of 2015), and Pubmed (from 1951 to April of 2015). A methodologic filter was applied to limit retrieval to clinical trials and the human population (detailed search strategy is provided in Supplemental Appendix 1) (10). We reviewed the reference lists of relevant articles and reviews as well as trials registered on the ClinicalTrials.gov website. We searched for conference proceedings from 2007 to 2014 using the Web of Science and abstracts presented at recent annual meetings of the American Society of Nephrology, the Canadian Society of Nephrology, the National Kidney Foundation, the European Renal Association-European Dialysis and Transplant Association, and the International Society of Nephrology.

Study Selection

We used the following eligibility criteria.

Studies. We included all randomized, controlled trials from January of 1946 to April of 2015. This included crossover, parallel arm, and cluster trials.

Participants. We included all adult patients (aged >18 years old) who received chronic hemodialysis in either an inpatient or outpatient setting, regardless of BP at baseline.

Intervention. We compared standard temperature dialysis with any method of cool dialysis. Different methods of cool dialysis include fixed reduction in dialysate temperature programmed patient cooling, isothermic dialysis, and negative energy balance using a biofeedback temperature monitoring device. We used the trials' definitions of standard and cool dialysis temperature. We only included bicarbonate-based dialysis and excluded acetate-based dialysis, because acetate-based dialysis is not used in common practice anymore. We did not include hemodiafiltration or acetate-free biofiltration in this review.

Outcomes Measures. We abstracted information on the following outcomes: mortality, hospitalization, quality of life (QOL), intradialytic hypotension, BP, cardiovascular events, access failure, system clotting, bleeding, dialysis adequacy, and symptoms, including discomfort caused by cold sensation and cramping. We did not assess the effect of cool dialysis on energy transfer, vasoactive peptides, or radiologic and diagnostic tests changes (e.g., changes in magnetic resonance, echocardiogram, or computed tomography).

Language. We included trials published in any language.

Publication Status. We reviewed all published and unpublished studies. Abstracts with relevant information were also reviewed.

Two investigators independently screened the search results for articles on the basis of the title or the title and abstract. The full-text article was retrieved for any citation considered potentially relevant by any investigator. Each of the investigators then independently assessed the eligibility of each article by using a pilot-tested, standardized form with written instructions. Any disagreement was resolved by consensus. If at least one of the characteristics was not met, the article was excluded. We used Cohen κ -value to measure agreement beyond chance between reviewers (11).

Data Extraction and Quality Assessment

We extracted data using a pilot-tested and standardized form. Two investigators independently extracted all relevant data from included trials. Results of data extraction were then compared, and any discrepancy was resolved by discussion. When the same results were presented in more than one publication, we included the publication with the most complete results. If results were incomplete or unclear, we contacted study authors for additional information.

We collected the following information from each trial: trial characteristics (author name, year of publication, country, language, number of centers, number of countries, and inclusion and exclusion criteria), patient characteristics (number, patients completing follow-up, age, comorbidities, and baseline BP), intervention and comparison characteristics (type of dialysate cooling, duration of intervention, and temperature of dialysate in each group), cointerventions (concomitant BP medication use), and outcomes. Also, we collected information about funding sources, conflict of interest statements, consent, and ethics approval.

Two reviewers independently assessed the risk of bias of the selected trials. To assess any risk of bias, different domains were considered for each outcome. These domains were adequate random sequence generation, concealment of allocation, adherence to the intention-to-treat principle, stopping early for benefit, proportion of patients lost to follow-up, selective reporting of outcomes, freedom from other biases, and blinding of patients, investigators, dialysis nurses, clinical outcome assessors, data collectors, and data analyzers.

To assess the confidence in the estimates of effect (*i.e.*, quality of evidence) across studies, we followed the Grading of Recommendations Assessment, Development and Evaluation GRADE approach (12) by making judgments about the risk of bias, publication bias, indirectness, imprecision, and inconsistency among different trials. To assess publication bias, we used the funnel plot and Egger linear regression test (13).

Data Synthesis and Analyses

The intervention effect estimates for each outcome were combined quantitatively (pooled) from different trials when appropriate using RevMan, version 5. When quantitative synthesis of the data was not appropriate, the results were summarized qualitatively. The Breslow–Day test was used to measure the percentage of total variation across studies caused by heterogeneity (I^2). Trial data were considered worthy of exploration of heterogeneity when the I^2 statistic was >50%. Attempts were also made to explain heterogeneity on the basis of the patient clinical characteristics and interventions of the trials included.

Rates were expressed as rate ratios and compared using generic inverse variance. Continuous data were expressed as mean differences. Overall results were pooled using a random effects model (14,15).

Sensitivity analyses to assess robustness of results and subgroup analyses to determine whether the summary effects vary in relation to clinical characteristics of the population in the included trials were prespecified. The treatment effects were examined according to risk of bias. Two subgroup analyses were undertaken. The first compared the effect of fixed temperature reduction compared

with biofeedback devices on different outcomes. The second compared the effect of cooling dialysate in patients with stable and unstable BPs at baseline.

Results

Study Selection

Study selection is presented in Figure 1; 4019 citations were screened, and a total of 26 trials were included in the review. The chance-corrected agreement for full-text eligibility was good ($\kappa=0.87$). Details about the excluded studies are presented in Supplemental Appendix 2. The most important reasons for exclusion included the use of acetate dialysis, duplicate data, and a lack of primary data.

Study Characteristics

Table 1 presents the details about characteristics of included trials (16–40) (B. Jo *et al.*, unpublished data). All

26 trials selected for the final review were crossover, randomized trials published in English. Nineteen trials had no dropouts. The other seven trials had patient dropout of varying degrees (8%–42%) (16–22). One trial was an international multicenter study (20) (27 centers and nine countries), and the rest were conducted in single centers.

Participants. The included trials involved 484 participants; 24 of 26 trials were small, with <20 participants (6–19 participants). The remaining two trials involved 95 and 128 participants (19,20). The main inclusion criteria were adults (≥ 18 years old) with variability in baseline BP and duration on hemodialysis.

Intervention. Overall, the duration of each trial was short. Eighteen trials reported the effect of cool dialysis on the basis of two sessions of hemodialysis (one session in each arm of the trial) (16,17,21,23,27–39,41). Three trials evaluated the intervention in three to six hemodialysis sessions (22–24). Only three trials followed patients longer




Figure 1. | Flow chart of the article selection process based on preferred reporting items for systematic reviews and meta-analyses.

Table 1. Characteristics of included trials

Study Country	Methods ^a	Participants	Intervention	Control	Outcomes	Exclusion Criteria
New Zealand (25)	Crossover design; allocation concealment: NR; sequence generation: NR; blinding: no; dropouts: 0	n=10; BP: five stable, five unstable; age 59.8±5.5 yr	Fixed temperature 35°C; duration: three sessions	Fixed temperature 36°C; duration: three sessions	Dialysis adequacy; BP; body temperature; symptoms; IDH	Circulation or vascular access problems, recent surgery, AKI, severe anemia, CAD
The Netherlands (16)	Crossover design; allocation concealment: NR; sequence generation: NR; blinding: NR; dropouts: 1	n=12 (eight men, four women); BP: all stable; age 69±6 yr	Fixed temperature 35.5°C; duration: one session	Fixed temperature 36.5°C; duration: one session	BP; body temperature; hemodynamics	CHF (NYHA≥3), severe CAD, DM
The Netherlands (26)	Crossover design; allocation concealment: NR; sequence generation: NR; blinding: NR; dropouts: 0	n=12 (seven men, five women); BP: all stable; age 64.7±2.6 yr	BTM (mean dialysate temperature 35.2°C); duration: one session	Fixed temperature 37.5°C; duration: one session	BP; body temperature; symptoms; IDH	CHF (EF≤25%), DM, CAD, NYHA≥3, using nitrates
United Kingdom (17)	Crossover design; allocation concealment: NR; sequence generation: NR; blinding: patients, not staff; dropouts: 1	n=10 (six men, four women); BP: not stable; age 67±2 yr	Fixed temperature 35°C; duration: one session	Fixed temperature 37°C; duration: one session	Dialysis adequacy; BP; body temperature; IDH; hemodynamics	Central venous catheter for access, severe CHF (NYHA≥3), heart transplant
United States (18)	Crossover design; allocation concealment: NR; sequence generation: NR; blinding: NR; dropouts: 8	n=19 (six men, five women); BP: not stable; age 67.5 yr	Fixed temperature 35.5°C; duration: nine sessions	Fixed temperature 37°C; duration: nine sessions	Dialysis adequacy; BP; symptoms; IDH	Symptoms not reported
United States (23)	Crossover design; allocation concealment: NR; sequence generation: NR; blinding: staff, not patients; dropouts: 0	n=10 (three men, seven women); BP: not stable; age 61.1±12.5 yr	Fixed temperature 35°C; duration: three sessions	Standard fixed temperature; duration: three sessions	Dialysis adequacy; BP; symptoms; IDH	Uncontrolled HTN, DM, unstable angina, noncompliance, frequent hospitalizations, variable weight gain

Table 1. (Continued)

Study Country	Methods ^a	Participants	Intervention	Control	Outcomes	Exclusion Criteria
Canada (19)	Crossover design; allocation concealment: NR; sequence generation: block center randomization; blinding: patients and nurses; dropouts: yes but no clear details; some patients studied more than once	<i>n</i> =128; BP: all stable; age 58 yr	Fixed temperature 35°C; duration: 13 sessions on average	Fixed temperature 37°C; duration: seven sessions on average	IDH	Pre-HD body temperature 36°C–36.5°C
Sweden (27)	Crossover design; allocation concealment: NR; sequence generation: NR; blinding: NR; dropouts: 0	<i>n</i> =10 (seven men, three women); BP: all stable; age 63 yr; time on HD: 3–49 mo	Fixed temperature 34.5°C; duration: one session	Fixed temperature 38.5°C; duration: one session	BP; body temperature	Not reported
United States (28)	Crossover design; allocation concealment: NR; sequence generation: NR; blinding: NR; dropouts: 0	<i>n</i> =13 (eight men, four women); BP: not stable; age 67.6±13.8 yr	Fixed temperature 35.5°C; duration: one session	Fixed temperature 37°C; duration: one session	BP; hemodynamics; complications	Central venous catheter for access, vascular access dysfunction, active medical condition
Japan (29)	Crossover design; allocation concealment: NR; sequence generation: NR; blinding: NR; dropouts: 0	<i>n</i> =7 (one man, six women); BP: not stable; age 53–75 yr	Fixed temperature 35.5°C; duration: one session	Fixed temperature 37°C; duration: one session	BP; symptoms	Not reported
United States (30)	Crossover design; allocation concealment: NR; sequence generation: NR; blinding: patients and investigators; dropouts: 0	<i>n</i> =12 (12 men); BP: not stable; age 62.5±3.6 yr; time on HD: 35.4±6.2 mo	Fixed temperature 35°C; duration: one session	Fixed temperature 37°C; duration: one session	BP; IDH; hemodynamics	Not reported
United States (31)	Crossover design; allocation concealment: NR; sequence generation: NR; blinding: NR; dropouts: 0	<i>n</i> =15; BP: not reported	BTM negative energy (dialysate temperature 35.7°C); duration: one session	BTM thermoneutral (dialysate temperature 37.1°C); duration: one session	Dialysis adequacy; BP; body temperature; IDH; hemodynamics; symptoms	Not reported

Table 1. (Continued)

Study Country	Methods ^a	Participants	Intervention	Control	Outcomes	Exclusion Criteria
Japan (32)	Crossover design; allocation concealment: NR; sequence generation: NR; blinding: NR; dropouts: 0	n=9; BP: not reported; age 43.5 yr	Fixed temperature 34°C; duration: one session	Fixed temperature 37°C; duration: one session	BP; hemodynamics	Not reported
Germany (33)	Crossover design; allocation concealment: NR; sequence generation: NR; blinding: NR; dropouts: 0	n=9; BP: not reported	BTM (program cooling by 0.2°C); duration: one session	Fixed temperature 37°C; duration: one session	Dialysis adequacy; BP; body temperature; hemodynamics	Not reported
United States (34)	Crossover design; allocation concealment: NR; sequence generation: NR; blinding: patients and providers; dropouts: 0	n=6; BP: not reported; age 55±11 yr	Fixed temperature 35°C; duration: one session	Fixed temperature 37°C; duration: one session	BP; body temperature	Not reported
Nine countries (20)	Crossover design (27 centers); allocation concealment: NR; sequence generation: central; blinding: no; dropouts: 21	n=95; BP: not stable; age 66±12 yr	BTM isothermic; duration: 12 sessions on average	BTM thermoneutral; duration: 12 sessions on average	Dialysis adequacy; BP; symptoms; IDH; complications	Recent surgery, severe anemia, vascular access problems, malignancy, ascitis, CHF (NYHA class 4), use of any antihypertensive medications
United States (41)	Crossover design; allocation concealment: NR; sequence generation: NR; blinding: single blinded; dropouts: 0	n=7 (three men, four women); BP: all stable; age 46.1±4.2 yr	Fixed temperature 35°C; duration: one session	Fixed temperature 37°C; duration: one session	BP; body temperature; sleep measures	Chronic infection, CHF, chronic lung disease, arthritis, organic brain disease, drug/alcohol abuse or past psychiatric disorders requiring treatment; taking β-adrenergic blockers, clonidine,

Table 1. (Continued)

Study	Country	Methods ^a	Participants	Intervention	Control	Outcomes	Exclusion Criteria
Australia (35)		Crossover design; allocation concealment: NR; sequence generation: NR; blinding: NR; dropouts: 0	n=8 (four men, four women); BP: not reported; age 54–79 yr	Fixed temperature 35.3°C±0.2°C; duration: one session	Fixed temperature 37.3°C±0.3°C; duration: one session	BP; body temperature	methyl dopa, antidepressants, sedatives, activating agents or pain medication, NSAIDs, and acetaminophen Not reported
United Kingdom (21)		Crossover design; allocation concealment: NR; sequence generation: NR; blinding: patients, not staff; dropouts: 1	n=10 (four men, six women); BP: not reported; age 65±11 yr	Fixed temperature 35°C; duration: one session	Fixed temperature 37°C; duration: one session	Dialysis adequacy; BP; body temperature; symptoms; IDH; QOL; hemodynamics	CHF (greater than NYHA class 3), cardiac transplant if it was not possible to obtain an echocardiogram Not reported
The Netherlands (36)		Crossover design; allocation concealment: NR; sequence generation: NR; blinding: NR; dropouts: 0	n=9 (four men, five women); BP: three of nine not stable; age 68.7±10.4 yr	Fixed temperature 35.5°C; duration: one session	Fixed temperature 37°C; duration: one session	BP; body temperature; IDH	
The Netherlands (37)		Crossover design; allocation concealment: NR; sequence generation: NR; blinding: NR; dropouts: 0	n=15 (seven men, eight women); BP: not reported; age 55 yr	Fixed temperature 35.5°C; duration: one session (1 h)	Fixed temperature 37.5°C; duration: one session (1 h)	BP; body temperature; symptoms; hemodynamics	Severe CAD, DM, CHF (LVEF<30%)
The Netherlands (38)		Crossover design; allocation concealment: NR; sequence generation: NR; blinding: NR; dropouts: 0	n=12 (seven men, five women); BP: all stable; age 56.67±15.95 yr	Fixed temperature 35.5°C; duration: one session	Fixed temperature 37.5°C; duration: one session	BP; body temperature	Not reported

Table 1. (Continued)

Study Country	Methods ^a	Participants	Intervention	Control	Outcomes	Exclusion Criteria
The Netherlands (39)	Crossover design; allocation concealment: NR; sequence generation: NR; blinding: NR; dropouts: 0	n=13; BP: all stable	BTM isothermic; duration: one session	BTM thermoneutral; duration: one session	Body temperature	Frequent IDH, severe CAD, severe CHF, central venous catheter for access
The Netherlands (22)	Crossover design; allocation concealment: NR; sequence generation: NR; blinding: NR; dropouts: 3	n=17; (eight men, six women); BP: not stable; age 60.9 ± 10.4 yr	BTM isothermic; BTM cooling 0.5°C below body temperature; duration: 1.5 sessions on average	BTM thermoneutral; duration: 1.5 sessions on average	BP; body temperature; symptoms; hemodynamics; IDH	Age >85 yr, not able to read and understand English, using central venous catheter for access
China (24)	Crossover design; allocation concealment: NR; sequence generation: NR; blinding: patients; dropouts: 0	n=9; BP: not reported; age 63 ± 2.2 yr	Fixed temperature 35°C; duration: two sessions	Fixed temperature 37.5°C; duration: two sessions	BP; body temperature; IDH; hemodynamics; dialysis adequacy	Not reported
Australia (40)	Crossover design; allocation concealment: NR; sequence generation: NR; blinding: NR; dropouts: 0	n=17 (nine men, eight women); BP: all stable; age 63.3 ± 3.2 yr	Fixed temperature 35°C; duration: NC	Fixed temperature 37°C; duration: NC	BP; Hemodynamics	Arthritis, severe PAD, α- or β-adrenergic blockers

NR, not reported; IDH, intradialytic hypotension; CAD, coronary artery disease; CHF, congestive heart failure; NYHA, New York Heart Association classification; DM, diabetes mellitus; BTM, biofeedback temperature monitoring; EF, ejection fraction; HTN, hypertension; HD, hemodialysis; NSAIDs, nonsteroidal anti-inflammatory drugs; QOL, quality of life; LVEF, left ventricular ejection fraction; NC, not clear; PAD, peripheral arterial disease.

^aNone of the studies stopped early for benefit. It is not clear if the studies had selective reporting of outcomes.




Figure 2. | Effect of low temperature dialysis on intradialytic hypotension. 95% CI, 95% confidence interval; BTM, biofeedback temperature monitoring.

than six sessions (18, 20, and 24 sessions on average, respectively) (18–20). Of note, trials with prolonged follow-up had a high dropout rate, and an intention-to-treat analysis was not followed. It is not clear if the dropouts were preferential in cool dialysis or not. Twenty trials used fixed temperature cooling of dialysate fluid (34°C–35.5°C). Six trials used a biofeedback temperature monitoring device

(one negative energy, two cooling, and three isothermic devices). In the standard dialysis group, the dialysate temperature ranged from 36°C to 37.5°C, except for in one trial that used a temperature of 38.5°C. Including or excluding this trial did not affect our results, and we decided to include it, although a temperature >37.5°C is higher than the standard and an unusual practice.




Figure 3. | Effect of low temperature dialysis on symptoms of discomfort. 95% CI, 95% confidence interval; BTM, biofeedback temperature monitoring.

Outcomes. No trial included mortality, hospitalization, cardiovascular events, access failure, bleeding, or system clotting as an outcome. One trial reported the effect of cooling dialysate on QOL; 12 trials reported intradialytic hypotension, 24 trials reported BP, nine trials reported dialysis adequacy, and ten trials reported symptoms of discomfort.

Risk of Bias within Studies

Seven trials described blinding of patients, four trials blinded health care providers or investigators, two trials reported no blinding, and blinding was unclear in the rest of the trials. Only one trial reported a central process for random sequence generation, and the rest did not report any methods for sequence generation or concealment of allocation. We encountered challenges when evaluating our criteria of selective reporting of outcomes, because published protocols for those trials were unavailable. None of the trials were stopped early for benefit or harm.

Intradialytic Hypotension

Thirteen trials reported intradialytic hypotension as an outcome. We summarize the definitions used by these trials to define intradialytic hypotension in Supplemental Appendix 3. Two trials (19,20) reported intradialytic hypotension as the proportion of sessions where it occurred. These studies (19,20) reported that 5.5%–25% of the dialysis sessions were complicated by intradialytic hypotension in the

cool dialysis group compared with 11.2%–50% of the sessions in the usual dialysate group. The pooled effect on the basis of 11 trials (Figure 2), which included a total of 552 hemodialysis sessions in 120 patients (with an average of 4.6 sessions per patient), showed that the rate of intradialytic hypotension was reduced by 70% (95% confidence interval [95% CI], 49% to 89%) with cool dialysis compared with standard dialysis ($I^2=0\%$).

Symptoms of Discomfort

The symptoms that were considered were discomfort on dialysis from feeling cold, shivering, or cramps. The reporting of symptoms of discomfort during dialysis was generally poor. Ten trials reported these symptoms, but they did not uniformly rate their severity. Cruz *et al.* (18) reported the proportion of sessions with symptoms on cold dialysate (13%). The pooled effect on the basis of nine trials (Figure 3), which included a total of 2548 hemodialysis sessions, showed that symptoms occur 2.95 (95% CI, 0.88 to 9.82) times more commonly with cool dialysis compared with standard dialysis ($I^2=0\%$).

Dialysis Adequacy

Nine trials reported dialysis adequacy as an outcome. Dheenan and Henrich (23) reported urea reduction rate. Levin and colleagues (22) reported no difference in urea kinetics, but no clear estimates were presented. The other seven trials reported Kt/V. Our results show that cool




Figure 4. | Effect of low temperature dialysis on dialysis adequacy. 95% CI, 95% confidence interval; BTM, biofeedback temperature monitoring.

dialysis did not have a significant effect on dialysis adequacy compared with standard dialysis, with a pooled Kt/V mean difference of -0.05 (95% CI, -0.09 to 0.01). We observed some heterogeneity in the results on this outcome across the trials ($I^2=63\%$) (Figure 4), which was partially explained ($I^2=18\%$) when we conducted a predefined sensitivity analysis on the basis of risk of bias.

BP

Twenty-four trials reported BP. Some reported a change in mean arterial pressure (MAP), intradialytic MAP, MAP posthemodialysis, changes in systolic and diastolic BP, or systolic and diastolic BP posthemodialysis. We choose to pool change in MAP (before and after hemodialysis), because it is less affected by baseline BP before hemodialysis and was one of the most frequently reported BP measures in the included trials. We meta-analyzed the results of ten trials that reported change in MAP (Figure 5). Our results show that cool dialysis is associated with significantly higher MAP of 12 mmHg (95% CI, 8 to 16 mmHg) compared with standard dialysis ($I^2=34\%$). Additionally, in Table 2, we summarize the effect on other measures of BP described in 14 trials that we were not able to mathematically pool.

QOL

The work by Selby *et al.* (21) was the only trial that reported the effect of cool dialysis on QOL. In this trial, there was no difference between cool and standard dialysis on QOL using the short form 36 health survey assessment tool. Ayoub and Finlayson (25), although not clearly assessing QOL, did indicate that 80% of the patients receiving cool dialysis self-reported a dramatic improvement in their general health.

Additional Analyses

According to our protocol for this systematic review, we explored the effects of cool dialysis on heterogeneous outcomes by subgroup analysis. We stratified by different interventions (fixed versus biofeedback temperature monitoring device cooling of dialysate) and patient characteristics (stability of baseline BP). The latter analysis was limited to the trials that reported baseline BP. The subgroup analyses did not explain heterogeneity and did not affect the overall pooled estimate of dialysis adequacy.

To assess the robustness of the pooled estimate, we conducted sensitivity analyses on the basis of risk of bias in studies. Excluding studies with significant loss to follow-up (18,20) resulted in partial explanation of heterogeneity ($I^2=63\%-18\%$) for dialysis adequacy without affecting the overall pooled estimates of the remaining trials.

Quality of Evidence for the Body of Evidence

Overall, the quality of evidence for the outcomes of interest across all studies was graded low or very low. Details about different domains assessing the quality of evidence using the Grading of Recommendations Assessment, Development and Evaluation approach are summarized in the evidence profile (Table 3). We did not find evidence for publication bias using funnel plots and Egger linear regression test (intradialytic hypotension [$P=0.41$], symptoms of discomfort [$P=0.13$], dialysis adequacy [$P=0.49$], and change in MAP [$P=0.17$]). We were not able to fit metaregression models to examine the association with loss to follow-up for dialysis adequacy because of insufficient data.

Discussion

Twenty-six crossover, randomized trials (including 484 patients) with varying degrees of methodologic rigor were




Figure 5. | Effect of low temperature dialysis on change in mean arterial pressure. 95% CI, 95% confidence interval; BTM, biofeedback temperature monitoring.

included in this systematic review. The results suggest that cool dialysis significantly reduces the rate of intradialytic hypotension. The available evidence suggests no negative effect on dialysis adequacy, with an increase in symptoms of discomfort of unclear severity.

This review has several strengths. This review extends the review by Selby and McIntyre (6) in many ways. We

have identified and analyzed four more trials in this review. The comprehensive and up-to-date search makes it unlikely that relevant trials were missed. All steps, including initial screening, trials selection, and data abstraction, were performed independently in duplicate to minimize any potential biases arising from subjectivity in these tasks. Additionally, we translated non-English

Table 2. Summary of BP measures in trials that did not report change in mean arterial pressure

Reference no.	Cool HD	Standard HD
25	Low BP group: intradialytic MAP: 92 ± 8.4 mmHg; postdialysis MAP: 95.9 ± 10.9 mmHg; stable BP group: intradialytic MAP: 99 ± 11.5 mmHg; postdialysis MAP: 102 ± 13.5 mmHg	Low BP group: intradialytic MAP: 81 ± 7.2 mmHg; $P < 0.01^a$; postdialysis MAP: 86.9 ± 7.3 mmHg; $P = 0.01^a$; stable BP group: intradialytic MAP: 96.5 ± 12.1 mmHg; postdialysis MAP: 100 ± 14.0 mmHg
16	Change in SBP: -6.0 ± 2 mmHg ^b ; change in DBP: -4.1 ± 7.6 mmHg ^b	Change in SBP: -0.8 ± 22.7 mmHg ^b ; change in DBP: -3.8 ± 12.4 mmHg ^b
26	SBP: 146 ± 5 mmHg before dialysis; 140 ± 6 mmHg after dialysis; DBP: 81 ± 3 mmHg before dialysis; 79 ± 4 mmHg after dialysis	SBP: 150 ± 5 mmHg before dialysis; 132 ± 4 mmHg after dialysis; DBP: 81 ± 2 mmHg before dialysis; 76 ± 3 mmHg after dialysis
17	SBP reduction: $2.71 \pm 0.97\%$; DBP reduction: $8.63 \pm 1.19\%$; intradialytic MAP reduction: $3.01 \pm 0.98\%$	SBP reduction: $-7.54 \pm 1.92\%$; $P < 0.001^a$; DBP reduction: $-4.99 \pm 1.4\%$; $P < 0.001^a$; intradialytic MAP reduction: $-8.99 \pm 1.71\%$; $P < 0.001$
18	Mean lowest intradialytic SBP: 102.5 ± 2.9 mmHg; mean lowest intradialytic DBP: 61.7 ± 2.3 mmHg; intradialytic MAP: 75.0 ± 2.3 mmHg; SBP: 132 ± 3.3 mmHg before dialysis; 118.1 ± 3.5 mmHg after dialysis; DBP: 73.7 ± 2.3 mmHg before dialysis; 69.2 ± 2.6 mmHg after dialysis	Mean lowest intradialytic SBP: 90.6 ± 2.5 mmHg; $P < 0.001^a$; mean lowest intradialytic DBP: 54.9 ± 2.2 mmHg; $P = 0.02^a$; intradialytic MAP: 66.8 ± 2.1 mmHg; $P = 0.002^a$; SBP: 132.7 ± 3.4 mmHg before dialysis; 109.0 ± 2.1 mmHg after dialysis; $P < 0.01^a$; DBP: 74.9 ± 3.0 mmHg before dialysis; 63.6 ± 1.9 mmHg after dialysis; $P = 0.01^a$
32	MAP did not change with time at the end of cool HD	MAP decreased by 10% at the end of normal HD; $P =$ significant
33	MAP increased in six of nine sessions ^b	MAP increased in one of nine sessions ^b
20	SBP reduction: -14.2 ± 16.5 mmHg; DBP reduction: -5.8 ± 8.1 mmHg	SBP reduction: -20.5 ± 15.7 mmHg; DBP reduction: -8.8 ± 8.4 mmHg; $P < 0.05^a$
41	Mean intradialytic SBP: 136.9 ± 11.4 mmHg; mean intradialytic DBP: 75.4 ± 6.0 mmHg; mean intradialytic MAP: 95.7 ± 7.5 mmHg	Mean intradialytic SBP: 130.7 ± 10.5 mmHg; mean intradialytic DBP: 72.3 ± 6.0 mmHg; average intradialytic MAP: 91.6 ± 7.4 mmHg
21	Mean SBP: 158.8 ± 14 mmHg; mean DBP: 78.6 ± 4 mmHg; average intradialytic MAP: 110.9 ± 7 mmHg	Mean SBP: 141.6 ± 17 mmHg; $P < 0.001^a$; mean DBP: 69.4 ± 5 mmHg; $P < 0.001^a$; average intradialytic MAP: 92.6 ± 10 mmHg; $P < 0.001^a$
36	Maximum decrease in intradialytic SBP: 21.8 ± 26.1 mmHg; maximum decrease in intradialytic DBP: -0.1 ± 19.2 mmHg; SBP: 130.3 ± 21.7 mmHg before dialysis; 131.6 ± 21.4 mmHg after dialysis; DBP: 72.2 ± 10.2 mmHg before dialysis; 63.7 ± 9.0 mmHg after dialysis	Maximum decrease in intradialytic SBP: 43.2 ± 20.6 mmHg; maximum decrease in intradialytic DBP: 14.9 ± 9.6 mmHg; SBP: 144.4 ± 25.9 mmHg before dialysis; 117.3 ± 26.1 mmHg after dialysis; DBP: 72.2 ± 10.2 mmHg before dialysis; 67.0 ± 10.9 mmHg after dialysis
37	SBP: 144 ± 24 mmHg before dialysis; 144 ± 26 mmHg after dialysis; DBP: 68 ± 14 mmHg before dialysis; 76 ± 12 mmHg after dialysis	SBP: 152 ± 22 mmHg before dialysis; 134 ± 23 mmHg after dialysis; DBP: 75 ± 9 mmHg before dialysis; 71 ± 13 mmHg after dialysis
22	SBP: 159 ± 35 mmHg before dialysis; 127 ± 39 mmHg after dialysis; $P < 0.05^c$; DBP: 82 ± 12 mmHg before dialysis; 70 ± 17 mmHg after dialysis; $P < 0.05^c$; intradialytic lowest SBP: 113 ± 30 mmHg	SBP: 151 ± 27 mmHg before dialysis; 122 ± 28 mmHg after dialysis; $P < 0.05^c$; DBP: 82 ± 11 mmHg before dialysis; 65 ± 12 mmHg after dialysis; $P < 0.05^c$; intradialytic lowest SBP: 104 ± 27 mmHg; $P = 0.08$
40	SBP: 127 ± 6.4 mmHg before dialysis; 134 ± 3.9 mmHg after dialysis; DBP: 76 ± 3.9 mmHg before dialysis; 81 ± 3.5 mmHg after dialysis	SBP: 126 ± 4.6 mmHg before dialysis; 127 ± 2.1 mmHg after dialysis; DBP: 74 ± 2.8 mmHg before dialysis; 74 ± 3.2 mmHg after dialysis

Effect estimates are presented as BP measure $\pm$ SD. HD, hemodialysis; MAP, mean arterial pressure; NS, not significant; SBP, systolic BP; DBP, diastolic BP.

^a P value comparing cool and standard HD.

^bAfter HD compared with before HD.

^c P value for end versus start of dialysis.

Table 3. Evidence profile

Quality Assessment	No. of Studies	Design	Limitations	Inconsistency	Indirectness	Imprecision	Other Considerations	No. of Hemodialysis Sessions and Patients		Effect: Absolute	Quality	Importance
								Cool Dialysis	Standard Dialysis			
Mortality—not reported	0	—	—	—	—	—	—	—	—	—	—	Critical
Quality of life (measured with SF-36; better indicated by higher values)	1	Randomized trial	Very serious ^a	No serious inconsistency	No serious indirectness	Very serious ^b	None	18 hemodialysis sessions in nine patients	18 hemodialysis sessions in nine patients	MD 1 higher (21.64 lower to 23.64 higher)	⊕○○○	Critical
Hospitalization—not reported	0	—	—	—	—	—	—	—	—	—	—	Critical
Intradialytic hypotension (measured with no. of events per dialysis session; rate ratio <1.0 implies that cool dialysis versus standard dialysis has a lower risk of the outcome)	11	Randomized trials	Very serious ^c	No serious inconsistency	No serious indirectness	No serious imprecision	None	264 hemodialysis sessions in 120 patients	264 hemodialysis sessions in 120 patients	Rate ratio 0.3 higher (0.11–0.51 higher)	⊕⊕○○	Critical
Symptoms of discomfort (cold or cramping; measured with no. of events per dialysis session; rate ratio <1.0 implies that cool dialysis versus standard dialysis has a lower risk of the outcome)	9	Randomized trials	Very serious ^d	No serious inconsistency	No serious indirectness	Serious ^e	None	1234 hemodialysis sessions in 141 patients	1234 hemodialysis sessions in 141 patients	Rate ratio 2.95 (0.88–9.82)	⊕○○○	Critical
Dialysis adequacy (measured with Kt/V; better indicated by lower values)	7	Randomized trials	Very serious ^f	No serious inconsistency ^g	No serious indirectness	No serious imprecision	None	2100 hemodialysis sessions in 137 patients	2100 hemodialysis sessions in 137 patients	MD 0.05 lower (0.09 lower to 0.01 higher)	⊕⊕○○	Important
Change in mean arterial pressure (measured with millimeters of mercury; better indicated by lower values)	10	Randomized trials	Very serious ^h	No serious inconsistency	No serious indirectness	No serious imprecision	None	731 hemodialysis sessions in 102 patients	731 hemodialysis sessions in 102 patients	MD 12 higher (8–16 higher)	⊕⊕○○	Important

Question: what are the benefits and harms of cool versus standard dialysis for patients with ESRD? Settings: chronic hemodialysis (inpatient or outpatient). — denotes no data available. SF-36, short form 36 health survey; MD, mean difference; ⊕○○○ very low, ⊕⊕○○ low.

^aUnclear method for sequence generation and concealment of allocation reported. Only patients were blinded; one of ten patients dropped out of the study (in the work by Selby *et al.* [21]).

^bOnly one trial reported quality of life. The 95% confidence interval of the results crossed zero, with a very wide 95% confidence interval ranging from considerable benefit to considerable harm.

^cUnclear methods of sequence generation and concealment of allocation in any of 11 studies; two of 11 studies reported patient blinding, one of 11 studies reported no blinding, and in seven of 11 studies, blinding was unclear.

^dUnclear methods of sequence generation and concealment of allocation in eight of nine studies; only one of nine studies reported a central process of sequence generation. Two of nine studies reported no blinding, one of nine studies reported no blinding, and in seven of 11 studies, blinding was unclear.

^eUnclear methods of sequence generation and concealment of allocation in six of seven studies; only one of seven studies reported central sequence generation. Three of seven studies reported blinding of patients only, two of seven studies clearly reported no blinding, and blinding was not clear in two of seven studies. Four of seven studies had patients drop out of the studies; in one study, 21 of 95 patients dropped out, and in another study, eight of 19 patients dropped out.

^fSubstantial heterogeneity ($I^2=63\%$) that was partially explained by excluding studies with loss to follow-up ($I^2=18\%$).

^gUnclear methods for sequence generation and concealment of allocation reported. Blinding: four of ten studies blinded patients, one of ten studies blinded staff, one of ten studies blinded providers, and one of ten studies blinded investigators.

articles. Finally, we analyzed sources of bias and explored reasons for diversity in the published literature.

This review has few limitations. The results of this review are inherently limited by the quality of the primary included studies. First, the majority of included studies are small with short follow-up times. Second, no studies reported long-term patient-important outcomes, which is a major limitation in this area. None of the included studies, except one, reported a central method of random sequence generation or allocation concealment. Also, none of the studies had appropriately blinded patients, investigators, dialysis nurses, clinical outcome assessors, data collectors, and data analyzers. Additionally, seven trials had significant patient dropout from 8% to 42%. One may assume that the high rates of dropouts in some studies may be because of intolerability. However, the lack of information about timing and reasons for the dropouts makes it difficult to know whether this assumption is true. Surprisingly, we did not observe any improvement in the methodologic quality of the trials in more recent reports compared with earlier ones. Also, we did not observe any progress in reporting long-term patient-important outcomes.

Despite our low confidence in the estimates of effect owing to imprecision and risk of bias, cool dialysis remains a promising therapy that, we believe, is not frequently used. This may be because of the lack of high-quality evidence to support its possible net benefit or physicians' beliefs about a negative side effect profile. Still, this intervention is quite simple and can be implemented in any dialysis center in the world without any additional cost other than training nursing staff on a new protocol.

Cool dialysis can be achieved by either a fixed reduction in the temperature of the dialysate fluid or use of a biofeedback device. It will be of interest to the nephrology community to have trials that directly compare the effect of fixed empirical reduction of dialysate temperature with isothermic or cool biofeedback temperature monitoring devices, because the former does not consider patients' variability in core temperature and access recirculation, but the latter does. This direct comparison may provide information about the optimal management for patients at high risk of intradialytic hypotension during hemodialysis without causing significant symptoms of discomfort.

Additionally, cool dialysis seems promising when evaluating other surrogate outcomes that we did not explore in this review. The study by Eldehni *et al.* (42) shows that hemodialysis results in a significant amount of magnetic resonance imaging changes signifying brain injury that could possibly be alleviated by the use of cold-temperature hemodialysis. Also, Parker *et al.* (41) have found that the use of cold-temperature hemodialysis may improve nocturnal sleep by decreasing sympathetic activation and sustaining the nocturnal skin temperature. The poor reporting of symptoms of discomfort and the lack of systematic evaluation of their severity continue to limit the ability to accurately assess the effect of cool dialysis on significant patient discomfort. This review highlights that high-quality, large, multicenter studies with long follow-up are needed to assess the effect of cool dialysis on long-term patient-important outcomes, like mortality, major adverse cardiovascular

events, hospitalization, patients' functional and cognitive status, and discomfort.

Acknowledgments

Part of this study was funded by the Ontario Strategy for Patient-Oriented Research Support Unit. A.X.G. is supported by the Adam Linton Chair in Kidney Health Analytics.

Disclosures

None.

References

- Grassmann A, Gioberge S, Moeller S, Brown G: ESRD patients in 2004: Global overview of patient numbers, treatment modalities and associated trends. *Nephrol Dial Transplant* 20: 2587–2593, 2005
- Daugirdas JT: Pathophysiology of dialysis hypotension: An update. *Am J Kidney Dis* 38(Suppl 4): S11–S17, 2001
- Rosa AA, Fryd DS, Kjellstrand CM: Dialysis symptoms and stabilization in long-term dialysis. Practical application of the CUSUM plot. *Arch Intern Med* 140: 804–807, 1980
- Zager PG, Nikolic J, Brown RH, Campbell MA, Hunt WC, Peterson D, Van Stone J, Levey A, Meyer KB, Klag MJ, Johnson HK, Clark E, Sadler JH, Teredesai P: “U” curve association of blood pressure and mortality in hemodialysis patients. Medical Directors of Dialysis Clinic, Inc. *Kidney Int* 54: 561–569, 1998
- Maggiore Q, Pizzarelli F, Zoccali C, Sica S, Nicolò F, Parlongo S: Effect of extracorporeal blood cooling on dialytic arterial hypotension. *Proc Eur Dial Transplant Assoc* 18: 597–602, 1981
- Selby NM, McIntyre CW: A systematic review of the clinical effects of reducing dialysate fluid temperature. *Nephrol Dial Transplant* 21: 1883–1898, 2006
- Altman DG, Schulz KF, Moher D, Egger M, Davidoff F, Elbourne D, Gøtzsche PC, Lang T; CONSORT GROUP (Consolidated Standards of Reporting Trials): The revised CONSORT statement for reporting randomized trials: Explanation and elaboration. *Ann Intern Med* 134: 663–694, 2001
- Moher D, Jones A, Lepage L; CONSORT Group (Consolidated Standards for Reporting of Trials): Use of the CONSORT statement and quality of reports of randomized trials: A comparative before-and-after evaluation. *JAMA* 285: 1992–1995, 2001
- Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JP, Clarke M, Devereaux PJ, Kleijnen J, Moher D: The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: Explanation and elaboration. *J Clin Epidemiol* 62: e1–e34, 2009
- Chapter 6: Searching for studies. Lefebvre C, Manheimer E, Glanville J: Cochrane Information Retrieval Methods Group: Chapter 6: Searching for studies. Available at: <http://www.cochrane-handbook.org/>. Accessed January 1, 2011.
- Cohen J: Weighted kappa: Nominal scale agreement with provision for scaled disagreement or partial credit. *Psychol Bull* 70: 213–220, 1968
- Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, Schünemann HJ; GRADE Working Group: GRADE: An emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 336: 924–926, 2008
- Egger M, Davey Smith G, Schneider M, Minder C: Bias in meta-analysis detected by a simple, graphical test. *BMJ* 315: 629–634, 1997
- Lau J, Ioannidis JP, Schmid CH: Summing up evidence: One answer is not always enough. *Lancet* 351: 123–127, 1998
- DerSimonian R, Laird N: Meta-analysis in clinical trials. *Control Clin Trials* 7: 177–188, 1986
- Beerenhout C, DeJagere T, van der Sande FM, Bekers O, Leunissen KM, Kooman JP: Haemodynamics and electrolyte balance: A comparison between on-line pre-dilution haemofiltration and haemodialysis. *Nephrol Dial Transplant* 19: 2354–2359, 2004
- Chesterton LJ, Selby NM, Burton JO, McIntyre CW: Cool dialysate reduces asymptomatic intradialytic hypotension and increases baroreflex variability. *Hemodial Int* 13: 189–196, 2009

18. Cruz DN, Mahnensmith RL, Brickel HM, Perazella MA: Midodrine and cool dialysate are effective therapies for symptomatic intradialytic hypotension. *Am J Kidney Dis* 33: 920–926, 1999
19. Fine A, Penner B: The protective effect of cool dialysate is dependent on patients' predialysis temperature. *Am J Kidney Dis* 28: 262–265, 1996
20. Maggiore Q, Pizzarelli F, Santoro A, Panzetta G, Bonforte G, Hannedouche T, Alvarez de Lara MA, Tsouras I, Loureiro A, Ponce P, Sulková S, Van Roost G, Brink H, Kwan JT; Study Group of Thermal Balance and Vascular Stability: The effects of control of thermal balance on vascular stability in hemodialysis patients: Results of the European randomized clinical trial. *Am J Kidney Dis* 40: 280–290, 2002
21. Selby NM, Burton JO, Chesterton LJ, McIntyre CW: Dialysis-induced regional left ventricular dysfunction is ameliorated by cooling the dialysate. *Clin J Am Soc Nephrol* 1: 1216–1225, 2006
22. van der Sande FM, Wystrychowski G, Kooman JP, Rosales L, Raimann J, Kotanko P, Carter M, Chan CT, Leunissen KM, Levin NW: Control of core temperature and blood pressure stability during hemodialysis. *Clin J Am Soc Nephrol* 4: 93–98, 2009
23. Dheenan S, Henrich WL: Preventing dialysis hypotension: A comparison of usual protective maneuvers. *Kidney Int* 59: 1175–1181, 2001
24. Yu AW, Ing TS, Zabaneh RI, Daugirdas JT: Effect of dialysate temperature on central hemodynamics and urea kinetics. *Kidney Int* 48: 237–243, 1995
25. Ayoub A, Finlayson M: Effect of cool temperature dialysate on the quality and patients' perception of haemodialysis. *Nephrol Dial Transplant* 19: 190–194, 2004
26. Beerenhout CH, Noris M, Kooman JP, Porrati F, Binda E, Morigi M, Bekers O, van der Sande FM, Todeschini M, Macconi D, Leunissen KM, Remuzzi G: Nitric oxide synthetic capacity in relation to dialysate temperature. *Blood Purif* 22: 203–209, 2004
27. Hegbrant J, Mårtensson L, Ekman R, Nielsen AL, Thysell H: Dialysis fluid temperature and vasoactive substances during routine hemodialysis. *ASAIO J* 40: M678–M682, 1994
28. Hoebe H, Abu-Alfa AK, Mahnensmith R, Perazella MA: Hemodynamics in patients with intradialytic hypotension treated with cool dialysate or midodrine. *Am J Kidney Dis* 39: 102–107, 2002
29. Jamil KM, Yokoyama K, Takemoto F, Hara S, Yamada A: Low temperature hemodialysis prevents hypotensive episodes by reducing nitric oxide synthesis. *Nephron* 84: 284–286, 2000
30. Jost CM, Agarwal R, Khair-el-Din T, Grayburn PA, Victor RG, Henrich WL: Effects of cooler temperature dialysate on hemodynamic stability in "problem" dialysis patients. *Kidney Int* 44: 606–612, 1993
31. Kaufman AM, Morris AT, Lavarias VA, Wang Y, Leung JF, Glabman MB, Yusuf SA, Levoci AL, Polaschegg HD, Levin NW: Effects of controlled blood cooling on hemodynamic stability and urea kinetics during high-efficiency hemodialysis. *J Am Soc Nephrol* 9: 877–883, 1998
32. Kishimoto T, Yamamoto T, Shimizu G, Horiuchi N, Hirata S, Nishitani H, Mizutani Y, Yamakawa M, Yamazaki Y, Maekawa M: Cardiovascular stability in low temperature dialysis. *Dial Transplant* 15: 329–333, 1986
33. Levin NW, Morris AT, Lavarias VA, Wang Y, Glabman MB, Leung JP, Yusuf SA, LeVoci AL, Polaschegg HD, Kaufman AM: Effects of body core temperature reduction on haemodynamic stability and haemodialysis efficacy at constant ultrafiltration. *Nephrol Dial Transplant* 11[Suppl 2]: 31–34, 1996
34. Levy FL, Grayburn PA, Foulks CJ, Brickner ME, Henrich WL: Improved left ventricular contractility with cool temperature hemodialysis. *Kidney Int* 41: 961–965, 1992
35. Schneditz D, Martin K, Krämer M, Kenner T, Skrabal F: Effect of controlled extracorporeal blood cooling on ultrafiltration-induced blood volume changes during hemodialysis. *J Am Soc Nephrol* 8: 956–964, 1997
36. van der Sande FM, Kooman JP, Burema JH, Hameleers P, Kerkhofs AM, Barendregt JM, Leunissen KM: Effect of dialysate temperature on energy balance during hemodialysis: Quantification of extracorporeal energy transfer. *Am J Kidney Dis* 33: 1115–1121, 1999
37. van der Sande FM, Gladziwa U, Kooman JP, Böcker G, Leunissen KM: Energy transfer is the single most important factor for the difference in vascular response between isolated ultrafiltration and hemodialysis. *J Am Soc Nephrol* 11: 1512–1517, 2000
38. van der Sande FM, Kooman JP, Konings CJ, Leunissen KM: Thermal effects and blood pressure response during postdilution hemodiafiltration and hemodialysis: The effect of amount of replacement fluid and dialysate temperature. *J Am Soc Nephrol* 12: 1916–1920, 2001
39. van der Sande FM, Rosales LM, Brenner Z, Kooman JP, Kuhlmann M, Handelsman G, Greenwood RN, Carter M, Schneditz D, Leunissen KM, Levin NW: Effect of ultrafiltration on thermal variables, skin temperature, skin blood flow, and energy expenditure during ultrapure hemodialysis. *J Am Soc Nephrol* 16: 1824–1831, 2005
40. Zitt E, Neyer U, Meusburger E, Tiefenthaler M, Kotanko P, Mayer G, Rosenkranz AR: Effect of dialysate temperature and diabetes on autonomic cardiovascular regulation during hemodialysis. *Kidney Blood Press Res* 31: 217–225, 2008
41. Parker KP, Bailey JL, Rye DB, Bliwise DL, Van Someren EJ: Lowering dialysate temperature improves sleep and alters nocturnal skin temperature in patients on chronic hemodialysis. *J Sleep Res* 16: 42–50, 2007
42. Eldehni MT, Odudu A, McIntyre CW: Randomized clinical trial of dialysate cooling and effects on brain white matter. *J Am Soc Nephrol* 26: 957–965, 2015
43. Dhondt A, Elout S, Verbeke F, Vanholder R: Dialysate and blood temperature during hemodialysis: Comparing isothermic dialysis with a single-pass batch system. *Artif Organs* 34: 1132–1137, 2010
44. Donauer J, Schweiger C, Rumberger B, Krumme B, Böhrer J: Reduction of hypotensive side effects during online-hemodiafiltration and low temperature haemodialysis. *Nephrol Dial Transplant* 18: 1616–1622, 2003
45. du Cheyron D, Terzi N, Seguin A, Valette X, Prevost F, Ramakers M, Daubin C, Charbonneau P, Parenti JJ: Use of online blood volume and blood temperature monitoring during haemodialysis in critically ill patients with acute kidney injury: A single-centre randomized controlled trial. *Nephrol Dial Transplant* 28: 430–437, 2013
46. Hecking M, Antlanger M, Winnicki W, Reiter T, Werzowa J, Haidinger M, Weichhart T, Polaschegg HD, Josten P, Exner I, Lorenz-Turnheim K, Eigner M, Paul G, Klausner-Braun R, Hörl WH, Sunder-Plassmann G, Säemann MD: Blood volume-monitored regulation of ultrafiltration in fluid-overloaded hemodialysis patients: Study protocol for a randomized controlled trial. *Trials* 13: 79, 2012
47. Odudu A, Eldehni MT, Fakis A, McIntyre CW: Rationale and design of a multi-centre randomised controlled trial of individualised cooled dialysate to prevent left ventricular systolic dysfunction in haemodialysis patients. *BMC Nephrol* 13: 45, 2012
48. Alappan R, Cruz D, Abu-Alfa AK, Mahnensmith R, Perazella MA: Treatment of severe intradialytic hypotension with the addition of high dialysate calcium concentration to midodrine and/or cool dialysate. *Am J Kidney Dis* 37: 294–299, 2001
49. Barendregt JN, Kooman JP, Buurma JH, Hameleers P, Kerkhofs AM, Leunissen KM, Van Der Sande FM: The effect of dialysate temperature on energy transfer during hemodialysis (HD). *Kidney Int* 55: 2598–2608, 1999
50. Bazzato G, Coli U, Landini S, Lucatello S, Fracasso A, Morachiello P, Righetto F, Scanferla F: Vascular stability and temperature monitoring in patients prone to dialysis-induced hypotension. *Contrib Nephrol* 41: 394–397, 1984
51. Bazzato G, Coli U, Landini S, Lucatello S, Fracasso A, Morachiello P, Righetto F, Scanferla F: Temperature monitoring in dialysis-induced hypotension. *Kidney Int Suppl* 17: S161–S165, 1985
52. Canaud B, Chenine L, Renaud S, Leray H: Optimal therapeutic conditions for online hemodiafiltration. *Contrib Nephrol* 168: 28–38, 2011
53. Coli U, Landini S, Lucatello S, Fracasso A, Morachiello P, Righetto F, Scanferla F, Onesti G, Bazzato G: Cold as cardiovascular stabilizing factor in hemodialysis: Hemodynamic evaluation. *Trans Am Soc Artif Intern Organs* 29: 71–75, 1983
54. Coli L, La Manna G, Comai G, Ursino M, Ricci D, Piccarri M, Locatelli F, Di Filippo S, Cristinelli L, Bacchi M, Balducci A, Aucella F, Panichi V, Ferrandello FP, Turchini R, Lambertini D,

- Mura C, Marinangeli G, Di Loreto E, Quarello F, Forneris G, Tancredi M, Morosetti M, Palombo G, Di Luca M, Martello M, Emiliani G, Bellazzi R, Stefoni S: Automatic adaptive system dialysis for hemodialysis-associated hypotension and intolerance: A noncontrolled multicenter trial. *Am J Kidney Dis* 58: 93–100, 2011
55. Ghasemi A, Shafiee M, Rowghani K: Stabilizing effects of cool dialysate temperature on hemodynamic parameters in diabetic patients undergoing hemodialysis. *Saudi J Kidney Dis Transpl* 19: 378–383, 2008
 56. Gotch FA, Keen ML, Yarian SR: An analysis of thermal regulation in hemodialysis with one and three compartment models. *ASAIO Trans* 35: 622–624, 1989
 57. Hsu HJ, Yen CH, Hsu KH, Lee CC, Chang SJ, Wu IW, Sun CY, Chou CC, Yu CC, Hsieh KF, Chen CY, Hsu CY, Weng CH, Tsai CJ, Wu MS: Association between cold dialysis and cardiovascular survival in hemodialysis patients. *Nephrol Dial Transplant* 27: 2457–2464, 2012
 58. Jefferies HJ, Burton JO, McIntyre CW: Individualised dialysate temperature improves intradialytic haemodynamics and abrogates haemodialysis-induced myocardial stunning, without compromising tolerability. *Blood Purif* 32: 63–68, 2011
 59. Keijman JM, van der Sande FM, Kooman JP, Leunissen KM: Thermal energy balance and body temperature: Comparison between isolated ultrafiltration and haemodialysis at different dialysate temperatures. *Nephrol Dial Transplant* 14: 2196–2200, 1999
 60. Kerr PG, van Bakel C, Dawborn JK: Assessment of the symptomatic benefit of cool dialysate. *Nephron* 52: 166–169, 1989
 61. Leunissen KM, Kooman JP, van Kuijk W, van der Sande F, Luik AJ, van Hooff JP: Preventing haemodynamic instability in patients at risk for intra-dialytic hypotension. *Nephrol Dial Transplant* 11 [Suppl 2]: 11–15, 1996
 62. Lima EQ, Silva RG, Donadi EL, Fernandes AB, Zanon JR, Pinto KR, Burdmann EA: Prevention of intradialytic hypotension in patients with acute kidney injury submitted to sustained low-efficiency dialysis. *Ren Fail* 34: 1238–1243, 2012
 63. Lindholm T, Thysell H, Yamamoto Y, Forsberg B, Gullberg CA: Temperature and vascular stability in hemodialysis. *Nephron* 39: 130–133, 1985
 64. Cadena M, Medel H, Rodriguez F, Flores P, Mariscal A, Franco M, Pérez-Grovas H, Escalante B: Isothermic vs thermoneutral hemodiafiltration evaluation by indirect calorimetry. Presented at the Annual International Conference of the IEEE Engineering in Medicine and Biology Society, Vancouver, British Columbia, Canada, August 21–24 2008
 65. Maggiore Q, Pizzarelli F, Sisca S, Catalano C, Delfino D: Vascular stability and heat in dialysis patients. *Contrib Nephrol* 41: 398–402, 1984
 66. Maggiore Q, Pizzarelli F, Zoccali C, Sisca S, Nicolò F: Influence of blood temperature on vascular stability during hemodialysis and isolated ultrafiltration. *Int J Artif Organs* 8: 175–178, 1985
 67. Maggiore Q: Isothermic dialysis for hypotension-prone patients. *Semin Dial* 15: 187–190, 2002
 68. Mahida BH, Dumler F, Zasuwa G, Fleig G, Levin NW: Effect of cooled dialysate on serum catecholamines and blood pressure stability. *Trans Am Soc Artif Intern Organs* 29: 384–389, 1983
 69. Marcen R, Orofino L, Quereda C, Pascual J, Ortuño J: Effects of cool dialysate in dialysis-related symptoms. *Nephron* 54: 356–357, 1990
 70. Marcén R, Quereda C, Orofino L, Lamas S, Teruel JL, Matesanz R, Ortuño J: Hemodialysis with low-temperature dialysate: A long-term experience. *Nephron* 49: 29–32, 1988
 71. Orofino L, Marcén R, Quereda C, Villafruela JJ, Sabater J, Matesanz R, Pascual J, Ortuño J: Epidemiology of symptomatic hypotension in hemodialysis: Is cool dialysate beneficial for all patients? *Am J Nephrol* 10: 177–180, 1990
 72. Panzetta G, Bianco F, Galli G, Ianche M, Savoldi S, Vianello S, Vidi E, Cicinato P: Thermal energy balance during hemodialysis: The role of the filter membrane. *G Ital Nefrol* 19: 425–431, 2002
 73. Parker KP, Bailey JL, Rye DB, Bliwise DL, Van Someren EJ: Insomnia on dialysis nights: The beneficial effects of cool dialysate. *J Nephrol* 21 [Suppl 13]: S71–S77, 2008
 74. Provenzano R, Sawaya B, Frinak S, Polaschegg HD, Roy T, Zasuwa G, Dumler F, Levin NW: The effect of cooled dialysate on thermal energy balance in hemodialysis patients. *ASAIO Trans* 34: 515–518, 1988
 75. Quereda C, Marcén R, Lamas S, Hernández-Jodra M, Orofino L, Sabater J, Villafruela J, Ortuño J: Hemodialysis associated hypotension and dialysate temperature. *Life Support Syst* 3 [Suppl 1]: 18–22, 1985
 76. Sherman RA, Faustino EF, Bernholz AS, Eisinger RP: Effect of variations in dialysate temperature on blood pressure during hemodialysis. *Am J Kidney Dis* 4: 66–68, 1984
 77. Sherman RA, Rubin MP, Cody RP, Eisinger RP: Amelioration of hemodialysis-associated hypotension by the use of cool dialysate. *Am J Kidney Dis* 5: 124–127, 1985
 78. van Kuijk WH, Hillion D, Savoie C, Leunissen KM: Critical role of the extracorporeal blood temperature in the hemodynamic response during hemofiltration. *J Am Soc Nephrol* 8: 949–955, 1997
 79. van Kuijk WH, Luik AJ, de Leeuw PW, van Hooff JP, Nieman FH, Habets HM, Leunissen KM: Vascular reactivity during haemodialysis and isolated ultrafiltration: Thermal influences. *Nephrol Dial Transplant* 10: 1852–1858, 1995
 80. van der Sande FM, Kooman JP, van Kuijk WH, Leunissen KM: Management of hypotension in dialysis patients: Role of dialysate temperature control. *Saudi J Kidney Dis Transpl* 12: 382–386, 2001
 81. Maggiore Q, Pizzarelli F, Dattolo P, Maggiore U, Cerrai T: Cardiovascular stability during haemodialysis, haemofiltration and haemodiafiltration. *Nephrol Dial Transplant* 15 [Suppl 1]: 68–73, 2000
 82. Brewster UC, Ciampi MA, Abu-Alfa AK, Perazella MA: Addition of sertraline to other therapies to reduce dialysis-associated hypotension. *Nephrology (Carlton)* 8: 296–301, 2003
 83. Maggiore Q, Pizzarelli F, Sisca S, Zoccali C, Parlongo S, Nicolò F, Creazzo G: Blood temperature and vascular stability during hemodialysis and hemofiltration. *Trans Am Soc Artif Intern Organs* 28: 523–527, 1982
 84. Maggiore Q, Dattolo P, Piacenti M, Morales MA, Pelosi G, Pizzarelli F, Cerrai T: Thermal balance and dialysis hypotension. *Int J Artif Organs* 18: 518–525, 1995
 85. Maggiore Q, Enia G, Catalano C, Misefari V, Mundo A: Effect of blood cooling on cuprophane-induced anaphylatoxin generation. *Kidney Int* 32: 908–911, 1987
 86. Nesrallah GE, Suri RS, Guyatt G, Mustafa RA, Walter SD, Lindsay RM, Akl EA: Biofeedback dialysis for hypotension and hypervolemia: A systematic review and meta-analysis. *Nephrol Dial Transplant* 28: 182–191, 2013
 87. Shahgholian N, Ghafourifard M, Shafiei F: The effect of sodium and ultra filtration profile combination and cold dialysate on hypotension during hemodialysis and its symptoms. *Iran J Nurs Midwifery Res* 16: 212–216, 2011
 88. Smirnov AV, Nesterova OB, Suglobova ED, Golubev RV, Vasil'ev AN, Vasil'eva IA, Verbitskaia EV, Korosteleva NI, Kostereva EM, Lebedeva EB, Levykina EN, Starosel'skiĭ KG, Lazeba VA: Clinical and laboratory evaluation of the efficiency of chronic hemodialysis treatment using acid succinate in patients with terminal renal failure. *Ter Arkh* 85: 69–75, 2013
 89. Teruel JL, Martins J, Merino JL, Fernández Lucas M, Rivera M, Marcén R, Quereda C, Ortuño J: Temperature dialysate and hemodialysis tolerance. *Nefrologia* 26: 461–468, 2006

Received: April 25, 2015 Accepted: November 9, 2015

Published online ahead of print. Publication date available at www.cjasn.org.

This article contains supplemental material online at <http://cjasn.asnjournals.org/lookup/suppl/doi:10.2215/CJN.04580415/-/DCSupplemental>.

Appendices:

Appendix 1: Search strategies

Ovid MEDLINE(R) 1946 to April Week 2 2015 -Search ran 15/04/2015

#	Searches	Search Type
1	randomized controlled trial.pt.	Advanced
2	controlled clinical trial.pt.	Advanced
3	randomized.ab.	Advanced
4	placebo.ab.	Advanced
5	randomly.ab.	Advanced
6	trial.ab.	Advanced
7	groups.ab.	Advanced
8	1 or 2 or 3 or 4 or 5 or 6 or 7 [mp=protocol supplementary concept, rare disease supplementary concept, title, original title, abstract, name of substance word, subject heading word, unique identifier]	Advanced
9	(animals not (humans and animals)).sh.	Advanced
10	8 not 9	Advanced
11	exp Renal Insufficiency/	Advanced
12	(kidney adj failure\$).mp. [mp=protocol supplementary concept, rare disease supplementary concept, title, original title, abstract, name of substance word, subject heading word, unique identifier]	Advanced
13	(kidney adj disease\$).mp. [mp=protocol supplementary concept, rare disease supplementary concept, title, original title, abstract, name of substance word, subject heading word, unique identifier]	Advanced
14	h?emodialysis.mp. [mp=protocol supplementary concept, rare disease supplementary concept, title, original title, abstract, name of substance word, subject heading word, unique identifier]	Advanced
15	exp Renal Dialysis/	Advanced
16	(renal adj dialysis).mp. [mp=protocol supplementary concept, rare disease supplementary concept, title, original title, abstract, name of substance word, subject heading word, unique identifier]	Advanced
17	(kidney adj dialysis).mp. [mp=protocol supplementary concept, rare disease supplementary concept, title, original title, abstract, name of substance word, subject heading word, unique identifier]	Advanced
18	(renal adj failure\$).mp. [mp=protocol supplementary concept, rare disease supplementary concept, title, original title, abstract, name of substance word, subject heading word, unique identifier]	Advanced

	identifier]	
19	(renal adj disease\$).mp. [mp=protocol supplementary concept, rare disease supplementary concept, title, original title, abstract, name of substance word, subject heading word, unique identifier]	Advanced
20	11 or 12 or 13 or 15 or 16 or 17 or 18 or 19	Advanced
21	exp Dialysis Solutions/	Advanced
22	dialysate\$.mp.	Advanced
23	(cool adj dialysate\$).mp. [mp=protocol supplementary concept, rare disease supplementary concept, title, original title, abstract, name of substance word, subject heading word, unique identifier]	Advanced
24	(cold adj dialysate\$).mp. [mp=protocol supplementary concept, rare disease supplementary concept, title, original title, abstract, name of substance word, subject heading word, unique identifier]	Advanced
25	(cool adj solution\$).mp. [mp=protocol supplementary concept, rare disease supplementary concept, title, original title, abstract, name of substance word, subject heading word, unique identifier]	Advanced
26	(cold adj solution\$).mp. [mp=protocol supplementary concept, rare disease supplementary concept, title, original title, abstract, name of substance word, subject heading word, unique identifier]	Advanced
27	21 or 22 or 23 or 24 or 25 or 26	Advanced
28	10 and 20 and 27	Advanced

Central through Wiley -Search Ran 15/04/2015 using title, abstract and keyword for the keyword searching

ID	Search
#1	MeSH descriptor Kidney Failure explode all trees
#2	kidney near disease*
#3	kidney near failure*
#4	MeSH descriptor Renal Replacement Therapy explode all trees
#5	kidney near dialysis
#6	renal near dialysis
#7	hemodialysis
#8	haemodialysis
#9	(#1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7 OR #8)

- #10 [MeSH descriptor Dialysis Solutions explode all trees](#)
- #11 [dialysis near solution*](#)
- #12 [dialysate*](#)
- #13 [cool near dialysate*](#)
- #14 [cold near dialysate*](#)
- #15 [cool near solution*](#)
- #16 [cold near solution*](#)
- #17 [\(#10 OR #11 OR #12 OR #13 OR #14 OR #15 OR #16\)](#)
- #18 [\(#9 AND #17\)](#)

PubMed -Search Ran 15/04/2015

Search	Most Recent Queries
#82	Search #57 and #72 and #81
#81	Search #74 or #75 or #76 or #77 or #78 or #80
#80	Search dialysis solutions [MeSH Terms]
#78	Search cold solution*
#77	Search cool solution*
#76	Search cold dialysate
#75	Search cool dialysate
#74	Search dialysate
#72	Search #59 or #60 or #61 or #62 or #63 or #64 or #65 or #66 or #67 or #70 or #71
#71	Search hemodialysis
#70	Search renal dialysis [MeSH Terms]
#67	Search end stage renal failure [tiab]
#66	Search end stage kidney failure [tiab]
#65	Search end stage kidney disease [tiab]
#64	Search end stage renal disease [tiab]
#63	Search end stage renal failure
#62	Search end stage kidney failure

- [#61](#) Search **end stage kidney disease**
- [#60](#) Search **end stage renal disease**
- [#59](#) Search **"kidney failure, chronic"[MeSH Terms]**
- [#57](#) Search **#55 NOT #56**
- [#56](#) Search **animals [mh] NOT humans [mh]**
- [#55](#) Search **#47 or #48 or #49 or #50 or #51 or #52 or #53 or #54**
- [#54](#) Search **groups [tiab]**
- [#53](#) Search **trial [tiab]**
- [#52](#) Search **randomly [tiab]**
- [#51](#) Search **drug therapy [sh]**
- [#50](#) Search **placebo [tiab]**
- [#49](#) Search **randomized [tiab]**
- [#48](#) Search **controlled clinical trial [pt]**
- [#47](#) Search **randomized controlled trial [pt]**

EMBASE (1947 to April Week 2 2015) - Search Ran 15/04/2015

#	Searches	Search Type
1	exp kidney failure/	Advanced
2	(renal adj disease*).mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer, drug manufacturer]	Advanced
3	(renal adj failure).mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer, drug manufacturer]	Advanced
4	exp DIALYSIS/	Advanced
5	exp HEMODIALYSIS/ or exp CONTINUOUS HEMODIALYSIS/	Advanced
6	(renal adj dialysis).mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer, drug manufacturer]	Advanced
7	(kidney adj dialysis).mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer, drug manufacturer]	Advanced
8	(kidney adj failure).mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer, drug manufacturer]	Advanced

9	(kidney adj disease\$).mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer, drug manufacturer]	Advanced
10	1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9	Advanced
11	exp dialysis fluid/	Advanced
12	exp DIALYSATE/	Advanced
13	(renal adj dialysis).mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer, drug manufacturer]	Advanced
14	(kidney adj dialysis).mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer, drug manufacturer]	Advanced
15	(cool adj dialysate).mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer, drug manufacturer]	Advanced
16	(cold adj dialysate).mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer, drug manufacturer]	Advanced
17	(cool adj solution).mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer, drug manufacturer]	Advanced
18	(cold adj solution).mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer, drug manufacturer]	Advanced
19	11 or 12 or 13 or 14 or 15 or 16 or 17 or 18	Advanced
20	randomized controlled trial.mp.	Advanced
21	exp controlled clinical trial/	Advanced
22	randomized.ab.	Advanced
23	placebo.ab.	Advanced
24	randomly.ab.	Advanced
25	trial.ab.	Advanced
26	groups.ab.	Advanced
27	20 or 21 or 22 or 23 or 24 or 25 or 26	Advanced
28	animals/ not humans.mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer, drug manufacturer]	Advanced
29	27 not 28	Advanced
30	10 and 19 and 29	Advanced

Appendix 2: Excluded studies

Study ID	Reason for exclusion
Alappan 2001 (49)	Additional Intervention – Ca concentration in dialysate adjusted
Barendregt 1999 (50)	Duplicate report
Bazzato 1984 (51)	Acetate dialysis
Bazzato 1985 (52)	Same data as previous citation
Canaud 2011 (53)	Used online HDF
Coli 1983 (54)	Same data as Bazzato et al
Coli 2011 (55)	No control group
Dhondt 2010 (28)	Low temp dialysis was not lower than 37 degrees
Donauer 2003 (29)	Used online hemodifiltration instead of conventional hemodialysis
Du Cheyron 2013 (30)	Patients had AKI not ESRD
Eldehni 2015 (26)	Did not report important patient outcomes
Ghasemi 2008 (56)	Acetate dialysis
Gotch 1989 (57)	Theoretical modeling for thermal balance
Hecking 2012 (31)	Included children and AKI
Hsu 2012 (58)	Observational study
Jefferies 2011 (59)	No control group, multi arm intervention study
Keijman 1999 (60)	Thermal changes only outcome measured
Kerr 1989 (61)	Acetate dialysis
Leunissen 1996 (62)	Review
Lima 2012 (63)	Patients with AKI
Lindholm 1985 (64)	Acetate dialysis
Cadena 2008 (65)	Thermal changes only outcome measured
Maggiore 1981 (5)	Acetate dialysis
Maggiore 1984 (66)	Review
Maggiore 1985 (67)	Acetate dialysis

Maggiore 2002 (68)	Review
Mahida 1983 (69)	Acetate dialysis
Marcen 1990 (70)	Acetate dialysis
Marcen 1988 (71)	Acetate dialysis
Orofino 1990 (72)	Acetate dialysis
Odudu 2012 (32)	Did not include patient important outcome
Panzetta 2002 (73)	Thermal changes only outcome measured
Parker 2008 (74)	Duplicate report
Provenzano 1988 (75)	Pilot study, thermal changes only outcome measured
Quereda 1985 (76)	Acetate dialysis
Sherman 1984 (77)	Acetate dialysis
Sherman 1985 (78)	Acetate dialysis
Van Kuijk 1997 (79)	Incomplete data
Van Kuijk 1995 (80)	Hemofiltration in intervention arm
Van Der Sande 2001 (81)	Review
Maggiore 2000 (82)	Comparing hemodialysis to hemofiltration
Brewster 2003 (83)	Additional intervention
Kerr 1989 (61)	Acetate dialysis
Maggiore 1982 (84)	Acetate dialysis
Maggiore 1995 (85)	Insufficient data
Maggiore 1987 (86)	Acetate dialysis
Nesrallah 2013 (87)	Review
Niyyar 2006 (88)	Duplicate Report
Shahgholian 2011 (89)	No control group
Smirnov 2013 (90)	Comparing succinate to bicarbonate dialysis solution
Teruel 2006 (91)	Not a randomized trial

Appendix 3: definitions of intra-dialytic hypotension by different studies

Article	IDH definition
Ayoub 2004 (33)	Systolic BP $\leq$ 90mm Hg accompanied by any of the following symptoms: nausea, vomiting, muscle cramps, dizziness, fainting
Beerenhout 2004a (34)	No definition provided
Chesterton 2009 (17)	SBP $\leq$ 100mmHg, even in the absence of symptoms, or a fall in SBP $>$ 10% of the predialysis reading in association with any of the classical symptoms of hypotension (e.g. headaches, cramps, lightheadedness)
Cruz 1999 (18)	Decrease in systolic blood pressure (SBP) of at least 20 mm Hg to a level $<$ 100 mm Hg at any time during HD
Dheenana 2001 (23)	An abrupt (over 10 to 15 min) decrement in blood pressure accompanied by symptoms or a decline of blood pressure requiring a nursing intervention
Fine 1996 (19)	Hypotensive symptoms occurring with a SBP below 100mmHg or with a 25% decrease in SBP in those patients with a basal SBP of 90-100mmHg or with a decrease of 50mmHg systolic
Jost 1993 (38)	Systolic blood pressure $<$ 90 mm Hg, the requirement for volume expansion therapy, or symptoms such as nausea, cramping, or vomiting
Kaufman 1998 (39)	Muscle cramps, nausea, vomiting, or episodes of hypotension requiring staff intervention were recorded
Maggiore 2002 (20)	Hypotensive episode was defined as the sudden onset of symptomatic hemodynamic changes that could be relieved by specific interventions. Specific interventions included Trendelenburg's position, administration of saline, administration of hypertonic solution, administration of plasma expander, and premature discontinuation of the dialysis session
Selby 2006 (21)	Systolic BP $\leq$ 100 mmHg, even in the absence of symptoms, or a fall in SBP $>$ 10% of the predialysis reading in association with any of the classical symptoms of hypotension (e.g. headaches, cramps, light-headedness)
Van Der Sande 1999 (44)	Systolic blood pressure $<$ 100 mm Hg, together with symptoms for which an infusion of saline 0.9% was necessary
Van Der Sande 2009 (22)	Show a difference of 20 mmHg in SBP between two different treatments
Yu 1995 (24)	Hypotension to a significant degree to warrant intervention