

Change from three times a week on-line hemodiafiltration to short daily on-line hemodiafiltration

FRANCISCO MADUELL, VICTOR NAVARRO, EDUARDO TORREGROSA, ASUNCIÓN RIUS, FERNANDO DICENTA, MARIA CARMEN CRUZ, and J. ANTONIO FERRERO

Department of Nephrology, Department of Cardiovascular Magnetic Resonance, and Department of Biochemistry, Hospital General de Castellón, Spain

Change from three times a week on-line hemodiafiltration to short daily on-line hemodiafiltration.

Background. Daily dialysis has shown excellent clinical results because a higher frequency of dialysis is more physiologic. On-line hemodiafiltration (OL-HDF) is a HDF technique that combines diffusion with high convection in which the dialysis fluid itself is used as a reinfusion solution. The aim of this study was to demonstrate the beneficial effect of the more effective dialysis schedule (daily dialysis) with the dialysis modality that offers the highest uremic toxin removal (on-line HDF).

Methods. Eight patients, six males and two females, on standard 4 to 5 hours three times a week OL-HDF (S-OL-HDF) were switched to daily OL-HDF (D-OL-HDF) 2 to 2½ hours six times per week. Dialysis parameters were identical during both periods and only frequency and dialysis time of each session were changed. Tolerance, uremic toxin removal, urea kinetics, biochemical and anemia profiles, blood pressure, and left ventricular hypertrophy were evaluated.

Results. D-OL-HDF was well accepted and tolerated. The disappearance of postdialysis fatigue was rapidly reported by patients. Patients maintained the same [time average concentration (TAC) and weekly single-pool Kt/V (spKt/V)] throughout the study. However, equivalent renal urea clearance (EKR), standard Kt/V and weekly urea reduction ratio (URR) were increased during D-OL-HDF. Weekly urea, creatinine, osteocalcin, β_2 -microglobulin, myoglobin, and prolactin reduction ratios were improved with D-OL-HDF. There was a significant decrease in predialysis plasma levels of urea, creatinine, acid uric, β_2 -microglobulin and homocysteine over 6 months. Phosphate binders were reduced and antihypertensive drugs were stopped. A 30% regression of left ventricular mass was observed.

Conclusion. The change from S-OL-HDF to D-OL-HDF was well tolerated. Disappearance of postdialysis fatigue, better dialysis adequacy, a higher removal of middle and large molecules, a reduction of phosphate binders, improvement of status nutritional, and an important reduction of cardiovascular risk factors were observed.

Key words: dialysis adequacy, daily hemodialysis, nutrition, on-line hemodiafiltration, solute removal.

Received for publication October 28, 2002

and in revised form January 8, 2003

Accepted for publication February 14, 2003

© 2003 by the International Society of Nephrology

The frequency of hemodialysis was established as thrice weekly in the 1960s and it has been mainly accepted and maintained for logistical, pragmatic, and economic reasons. However, there is a growing interest in the use of more frequent dialysis schedules since long-term experiences using higher frequencies have shown good results [1–5]. The first experiences with daily dialysis were reported in 1967 [6] and two predominant treatment schedules have been established. Short daily dialysis [1, 2, 4] has been used for more than 20 years and slow nightly dialysis [5, 7, 8] has been used since 1995. Although there are only a few studies and the number of patients is limited, they show excellent clinical results because a higher frequency of dialysis is more physiologic and it decreases fluctuations in fluid volume, solutes, and electrolytes. Improvements in comfort during and between dialysis sessions, clinical and biochemical parameters, anemia correction, hypertension control, nutrition status, and quality of life have been reported.

On-line hemodiafiltration (OL-HDF) could offer an optimal form of extracorporeal treatment for dialysis patients. This technique, which combines diffusion with high convection [infusion flux (Qi) 50 to 200 mL/minute or 3–12 L/hour], provides the highest clearances per unit of surface area for small, medium, and large-sized molecules. The same dialysis fluid, free of toxins and pyrogens, is used as substitution solution. It is a safe, well-tolerated technique, and good clinical results have been observed [9–11].

In this study, we sought to combine the more physiologic and effective dialysis schedule (daily dialysis) with the dialysis modality that offers higher solute and uremic toxin removal (OL-HDF). The aim of this study was to change patients from standard 4 to 5 hours three times a week OL-HDF to 2 to 2½ hours six times a week OL-HDF, and evaluate the impact of higher frequency therapy on solute removal capacity and on biochemical and clinical outcomes.

METHODS

This was a single-center, prospective, and nonrandomized study. Eight patients, six males and two females,

with a mean age of 65.9 ± 14 years (range, 41 to 80 years), stable on hemodialysis over a period of 68.4 ± 43 months and on standard 4 to 5 hours three times a week OL-HDF (S-OL-HDF) during the last 36.3 ± 21 months (range, 8 to 68 months) were switched to daily OL-HDF (D-OL-HDF) 2 to 2½ hours six times per week. The underlying renal diseases were chronic glomerulonephritis in two patients, nephroangiosclerosis in two patients, polycystic kidney disease in one patient, chronic tubulointerstitial nephritis in one patient, diabetic nephropathy in one patient, and undiagnosed nephropathy in one patient. All patients signed consent forms approved by the Hospital Research Committee.

The S-OL-HDF parameters were as follows: dialysis buffer with bicarbonate, 1.8 m² high-flux polysulfone filter (HF80, Fresenius, Bad Homburg, Germany), blood flow (Qb) 445 ± 54 mL/min (range, 350 to 560 mL/min), dialysate flow (Qd) 800 mL/min minus the infusion flow (Qi), which ranged from 80 to 150 mL/min and a Fresenius 4008 dialysis monitor. Reinfusion was always performed in a postdilutional mode. On D-OL-HDF, these parameters were the same and only the frequency and dialysis duration of each session were changed. All patients had native arteriovenous fistulas. The dialysers were not reused. Residual renal function was negligible in all patients. Before the study period, six patients used 15-gauge needles and two used 14-gauge needles. All patients used 15-gauge needles while on D-OL-HDF (Qb was limited to 500 mL/min).

The following urea kinetic parameters were calculated: single-pool second-generation Daugirdas Kt/V (spKt/V) [12], equilibrated Kt/V (eKt/V) [13], urea reduction ratio (URR), time average concentration (TAC), and normalized protein catabolic rate (nPCR). Moreover, different yardsticks proposed to measure the dialysis dose delivered by the different dialysis regimens have also been included: weekly spKt/V, weekly eKt/V, the equivalent renal urea clearance (EKR) introduced by Casino and Lopez [14], the standard Kt/V described by Gotch [15] and, finally, we calculated the weekly URR and the time average deviation (TAD) introduced by Lopot and Válek [16].

To evaluate removal capacity between both treatments modes, "in vivo" removal of a wide spectrum of solutes was studied. Pre- and postdialysis concentrations of urea (60 D), creatinine (113 D), osteocalcin (5800 D), β_2 -microglobulin (11,800 D), myoglobin (17,000 D), and prolactin (23,000 D) were measured. Pretreatment blood samples were drawn immediately after access needle insertion. Posttreatment samples were drawn from the arterial blood line 60 seconds after decreasing the blood flow rate to 50 mL/min. Osteocalcin concentrations were measured by a commercial immunometric assay, Immulite (DPC Diagnostics, Los Angeles, CA, USA) osteocalcin. The normal range of osteocalcin values is 3.1 to 13.7

ng/mL in our laboratory. Serum β_2 -microglobulin was measured using the Quantex β_2 -microglobulin commercial immunoturbidimetry, which has a normal range of 1.1 to 2.4 mg/L. Myoglobin concentrations were measured by a commercial immunoenzymatic "sandwich" assay (Access; Beckman, Fullerton, CA, USA), which has a normal range of 0 to 70 ng/mL. Prolactin concentrations were measured by a commercial immunometric assay, Immulite 2000 DPC, which has a normal range of 3 to 30 ng/mL. The pre- to posttreatment reduction ratios in plasma for osteocalcin, β_2 -microglobulin, myoglobin, and prolactin were determined after correcting postdialysis concentrations using the method of Bergström and Wehle [17].

The second part of study was designed to evaluate biochemical and clinical outcomes. The last 6 months on S-OL-HDF, during which conditions of treatment were not varied, were taken as the baseline period, before switching the patients to D-OL-HDF. Treatment was carried out over 6 months. Each month at mid-week, predialysis plasma analyses for hemoglobin, hematocrit, ferritin, transferrin saturation, fibrinogen, urea, creatinine, sodium, potassium, uric acid, bicarbonate, calcium, phosphorus, intact parathyroid hormone (iPTH), serum protein, albumin, prealbumin, transferrin, total cholesterol, high-density lipoprotein (HDL) cholesterol, low-density lipoprotein (LDL) cholesterol, triglycerides, β_2 -microglobulin, myoglobin, and C-reactive protein were carried out, as well as a quarterly analysis of homocysteine. Homocysteine was determined by the IMx method (IMX System; Abbott, Wiesbaden, Germany), using fluorescence polarization immunoassay technology.

At baseline and each month, each patient completed a fatigue index questionnaire concerning the intensity and duration of frequency of postdialysis fatigue [18]. Fatigue intensity was scored as follows: 0, none; 1, mild (noticeable but without effect); 2, moderate (felt sluggish); 3, severe (required rest); or 4, overwhelming (slept). Fatigue duration was scored as follows: 0, none; 1, mild (one to 4 hours); 2, moderate (6 to 12 hours); 3, severe (12 to 24 hours); or 4, overwhelming (more than 24 hours).

Predialysis blood pressure was measured immediately before treatment with an automatic blood pressure monitor (BPM). Left ventricular hypertrophy (LVH) was evaluated by magnetic resonance imaging (MRI). Measurements included left ventricular internal diameter at end-diastole (LVIDD) and left ventricular internal diameter at end-systole (LVIDS), interventricular septal thickness (IVS), anterior wall thickness at end-diastole (AWT), posterior wall thickness at end-diastole (PWT), lateral wall thickness at end-diastole (LWT), inferior wall thickness at end-diastole (IWT), left atrium dimension (LA) and aortic root dimension (AO). Left ventricular wall mass (LVM) was calculated by outlining endo- and epicardial contours on a stack of image sections of a specific

Table 1. Change from three times a week on-line hemodiafiltration (OL-HDF) to short daily on-line hemodiafiltration (D-OL-HDF): Comparison of urea kinetics during the two study periods

	Baseline	Month 3	Month 6
spKt/V	2.30 ± 0.20	1.13 ± 0.15 ^b	1.11 ± 0.11 ^b
eKt/V	1.96 ± 0.17	0.90 ± 0.12 ^b	0.88 ± 0.08 ^b
URR %	84.3 ± 2.5	64.2 ± 5.3 ^b	63.3 ± 4.2 ^b
Weekly spKt/V	6.90 ± 0.59	6.78 ± 0.91	6.67 ± 0.64
Weekly eKt/V	5.88 ± 0.52	5.39 ± 0.75 ^a	5.30 ± 0.50 ^a
EKR mL/min	19.2 ± 0.5	24.2 ± 2.6 ^b	23.8 ± 1.9 ^b
stdKt/V	2.62 ± 0.1	3.87 ± 0.3 ^b	3.86 ± 0.2 ^b
Weekly URR %	253 ± 7.5	385 ± 32 ^b	380 ± 25 ^b
TAC _{BUN} mg/dL	27.3 ± 7.1	26.7 ± 8.6	27.7 ± 6.1
TAD _{BUN} mg/dL	9.69 ± 2.3	6.24 ± 2.1 ^b	6.26 ± 1.3 ^b

Abbreviations are: URR, urea reduction ratio; spKt/V, single-pool Kt/V; eKt/V, equilibrated Kt/V; stdKt/V, standard Kt/V; TAC, time average concentration; TAD, time average deviation; ERK, equivalent renal urea clearance.

^a $P < 0.05$; ^b $P < 0.01$ with respect to baseline value (ANOVA repeated measures)


time frame encompassing the ventricle, using Simpson's rule algorithm [19]. LVM index (LVMI) was calculated by dividing LVM by body surface area. Left ventricular end-diastolic volume (LVEDV), end-systolic volume (LVESV), and ejection fraction (EF) were also calculated. All studies were performed and analyzed by the same experienced cardiologist, who was blinded to the patients' data. Studies were performed on a mid-week nondialysis day in S-OL-HDF and on a mid-week day before dialysis in D-OL-HDF. Specific criteria were used to determine the presence of LVH (men, LVM >158 g or LVMI >83 g/m²; women, LVM >123 g or LVMI >73 g/m²) [19].

The results were expressed as the arithmetic mean ± standard deviation. Each patient served as his/her own control. The Student *t* test (paired data) and analysis of variance (ANOVA) test (repetitive data) were used in the analysis of differences in quantitative variables. A value of $P < 0.05$ was considered statistically significant.

RESULTS

The duration per dialysis session changed from 274 ± 25 minutes (range, 240 to 300 minutes) in S-OL-HDF to 133 ± 12 minutes (range, 120 to 150 minutes) in D-OL-HDF ($P < 0.01$) but weekly dialysis time was similar, 822 minutes versus 798 minutes (NS), respectively. Q_b was 445 ± 59 mL/min in S-OL-HDF versus 438 ± 38 mL/min in D-OL-HDF (NS). Mean session value for reinfusion volume in S-OL-HDF was 30.7 ± 6 L and 14.6 ± 2 L in D-OL-HDF ($P < 0.01$), and weekly reinfusion volume was 95.8 ± 21 L and 88.1 ± 9 L (NS), respectively.

Daily OL-HDF was well accepted and tolerated by all patients. No local infections, thromboses, or bleeding of the vascular access occurred. The number of nausea, dizziness, cramps, and hypotensive episodes did not change between the two treatment schedules. In the first

**Fig. 1.** Percentage variation of different dialysis dose evaluations with change from three times a week on-line hemodiafiltration (OL-HDF) to short daily on-line hemodiafiltration (D-OL-HDF). Abbreviations are: spKt/V, single-pool Kt/V; eKt/V, equilibrated Kt/V; ERK, equivalent renal urea clearance; stdKt/V, standard Kt/V; URR, urea reduction ratio.

4 weeks, the patients reported a rapid improvement in the following symptoms: headaches in three patients, sleep disorders in three patients, sexual disorders in two patients, thoracic pain in one patient, increased appetite in five patients, and decreased thirst in two patients. The most significant benefit was the disappearance of postdialysis fatigue. Fatigue intensity scores diminished from 1.88 ± 1.2 in S-OL-HDF to 0.38 ± 0.7 in S-OL-HDF ($P < 0.01$), and fatigue duration scores decreased from 1.75 ± 1.4 in S-OL-HDF to 0.25 ± 0.5 in S-OL-HDF ($P < 0.01$).

Urea kinetics

Throughout the study, patients maintained the same TAC, weekly spKt/V, and weekly eKt/V (Table 1). However, EKR, standard Kt/V, and weekly URR were significantly increased in D-OL-HDF. The percentage variation of D-OL-HDF dialysis dose, measured by different methods, is shown in Figure 1. The outcome of pre- and postdialysis blood urea nitrogen (BUN), TAC_{BUN}, and TAD_{BUN} is shown in Figure 2.

Solute removal

Mean session values for urea, creatinine, osteocalcin, β_2 -microglobulin, myoglobin, and prolactin reduction ratios were lower on D-OL-HDF; however, the weekly reduction ratios were significantly increased on D-OL-HDF (Table 2). Figure 3 shows the weekly percentage increase in the efficacy of D-OL-HDF over a broad spectrum of solutes in comparison with S-OL-HDF.

There was a significant decrease in predialysis levels of urea, creatinine, uric acid (Table 3), and large molecules (β_2 -microglobulin and myoglobin) over 6 months (Fig. 4). The biochemical outcomes for predialysis so-




Fig. 2. Evolution of pre and postdialysis blood urea nitrogen (BUN), time average concentration (TAC)_{BUN} and time average deviation (TAD)_{BUN} switching from three times a week on-line hemodiafiltration (OL-HDF) to short daily on-line hemodiafiltration (D-OL-HDF).




Fig. 3. Increase in the weekly percentage removal of a broad spectrum of solutes with short daily on-line hemodiafiltration (D-OL-HDF) in comparison with three times a week on-line hemodiafiltration (OL-HDF). Abbreviations are: Crea, creatinine; Osteo, osteopontin; β_2 -m, microglobulin; Myo, myoglobin; PRL, prolactin.

Table 2. Comparison of urea, creatinine, osteocalcin, β_2 -microglobulin, myoglobin, and prolactin reduction ratios between three times a week standard on-line hemodiafiltration (S-OL-HDF) and short daily on-line hemodiafiltration (D-OL-HDF)

Reduction ratio	S-OL-HDF	D-OL-HDF	<i>P</i> value
Session			
Urea %	84.3 $\pm$ 3	64.2 $\pm$ 5	<0.001
Creatinine %	73.1 $\pm$ 2	56.7 $\pm$ 3	<0.001
Osteocalcin %	71.3 $\pm$ 7	60.6 $\pm$ 7	<0.001
β_2 -microglobulin %	79.8 $\pm$ 4	66.3 $\pm$ 11	<0.05
Myoglobin %	65.7 $\pm$ 5	56.5 $\pm$ 10	<0.05
Prolactin %	65.6 $\pm$ 4	57.1 $\pm$ 9	<0.05
Weekly			
Urea %	253 $\pm$ 8	385 $\pm$ 32	<0.001
Creatinine %	219 $\pm$ 5	340 $\pm$ 15	<0.001
Osteocalcin %	214 $\pm$ 21	363 $\pm$ 39	<0.001
β_2 -microglobulin %	239 $\pm$ 11	398 $\pm$ 66	<0.001
Myoglobin %	197 $\pm$ 14	339 $\pm$ 62	<0.001
Prolactin %	196 $\pm$ 13	343 $\pm$ 55	<0.001

dium, potassium, bicarbonate, calcium, phosphorus, and iPTH are shown in Table 3. Although serum phosphate did not change, phosphate binders (calcium carbonate in seven patients and calcium acetate plus aluminium hydroxide in one patient) were reduced from 7.3 ± 3 tablets/day on S-OL-HDF to 2.9 ± 3 tablets/day after 3 months and 2.85 ± 4 after 6 months ($P < 0.001$) on D-OL-HDF. The mean baseline value for β_2 -microglobulin (29.5 ± 2 mg/L) was 15 times higher than normal range and myoglobin (214 ± 75 mg/L) was three times higher than maximal normal range. The mean baseline predialysis level of osteocalcin (18.4 ± 19 mg/L) was slightly higher than maximal normal range, and the baseline predialysis levels of prolactin (16.4 ± 10 ng/mL) were within the normal range and remained constant during the 6-month study period (Table 3). In the diabetic patient, the insulin

dose was reduced from 24 IU/day at baseline to 12 IU/day after 2 months on D-OL-HDF.

Hematologic monitoring

Most hematologic parameters in our study did not show any significant changes (Table 4). Only the predialysis ferritin decreased from 473 ± 263 ng/mL at baseline to 312 ± 147 ng/mL after 3 months ($P < 0.01$) and to 290 ± 177 ng/mL after 6 months ($P < 0.01$). Erythropoietin doses (3250 ± 2052 IU/week) were not varied during the study and intravenous iron was increased from 24 to 46 mg/week (sodium ferric gluconate complex) during the study period.

Nutrition and inflammation parameters

Interdialytic weight gain per session decreased from 2.17 ± 1.5 kg to 1.25 ± 0.5 kg, but weekly interdialytic weight gain rose from 7.62 ± 1.5 kg on S-OL-HDF to 8.48 ± 2.5 kg on D-OL-HDF ($P < 0.001$) (Table 5). Mean nPCR increased from 0.93 ± 0.2 g/kg/day on S-OL-HDF to 1.18 ± 0.3 g/kg/day after 3 months ($P < 0.05$) and 1.13 ± 0.2 g/kg/day after 6 months on D-OL-HDF. There were no significant changes in serum protein, albumin, prealbumin, transferrin, total cholesterol, HDL cholesterol, LDL cholesterol, triglycerides, fibrinogen and C-reactive protein. Body weight (measured as dry weight after dialysis) increased from 67.8 ± 8 kg on S-OL-HDF to 68.3 ± 8 kg after 3 months and 69.4 ± 8 kg after 6 months (NS) on D-OL-HDF.

Blood pressure monitoring

During the baseline period, three patients had hypertension (predialysis mean blood pressure >110 mm Hg)

Table 3. Change from three times a week on-line hemodiafiltration (OL-HDF) to short daily on-line hemodiafiltration (D-OL-HDF). Predialysis levels of sodium, potassium, bicarbonate, urea, creatinine, uric acid, calcium, phosphorus, intact PTH (iPTH), osteocalcin and prolactin

	Baseline	Month 1	Month 2	Month 3	Month 4	Month 5	Month 6
Sodium <i>mmol/L</i>	139 ± 1	140 ± 4	140 ± 4	139 ± 4	140 ± 5	141 ± 1	139 ± 4
Potassium <i>mmol/L</i>	5.1 ± 0.5	4.5 ± 0.4	4.6 ± 0.7	4.7 ± 0.5	4.6 ± 0.7	4.7 ± 0.8	4.8 ± 0.5
Bicarbonate <i>mmol/L</i>	22.0 ± 2	23.6 ± 3	23.3 ± 2	24.4 ± 3 ^a	23.5 ± 3	25.6 ± 3 ^b	24.5 ± 3 ^a
Uric acid <i>mg/dL</i>	6.08 ± 1	5.12 ± 1 ^b	4.73 ± 1 ^b	5.03 ± 1 ^b	4.80 ± 1 ^b	4.64 ± 1 ^b	4.71 ± 1 ^b
Urea <i>mg/dL</i>	105 ± 20	83 ± 18 ^b	83 ± 17 ^b	83 ± 24 ^b	91 ± 10 ^a	82 ± 15 ^b	87 ± 15 ^b
Creatinine <i>mg/dL</i>	7.38 ± 1	6.19 ± 1 ^b	5.76 ± 1 ^b	6.03 ± 1 ^b	5.96 ± 1 ^b	6.23 ± 1 ^b	6.34 ± 1 ^b
Calcium <i>mg/dL</i>	9.8 ± 0.3	9.7 ± 0.7	9.6 ± 0.7	9.6 ± 0.6	9.6 ± 0.4	9.5 ± 0.5	9.8 ± 0.6
Phosphorus <i>mg/dL</i>	4.29 ± 1	4.54 ± 1	4.68 ± 1	4.11 ± 1	4.81 ± 1	4.46 ± 1	4.65 ± 1
iPTH <i>pg/mL</i>	136 ± 111	275 ± 288	176 ± 156	171 ± 157	189 ± 171	161 ± 141	117 ± 90
Osteocalcin <i>mg/L</i>	18.4 ± 19	19.6 ± 19	19.0 ± 20	23.0 ± 18	22.1 ± 17	19.8 ± 16	17.4 ± 14
Prolactin <i>ng/mL</i>	16.4 ± 10	17.3 ± 11	19.4 ± 14	18.9 ± 12	17.8 ± 12	19.8 ± 13	17.0 ± 12

^a $P < 0.05$. ^b $P < 0.01$ with respect to baseline value (ANOVA repeated measures)




Fig. 4. Change from three times a week on-line hemodiafiltration (HDF) to short daily on-line hemodiafiltration (D-OL-HDF). (A) Pre-dialysis serum β_2 -microglobulin (11,800 D) over 6 months. (B) Myoglobin (17,200 D) levels over 6 months. $P < 0.01$, analysis of variance (ANOVA) repeated measures.

and two were receiving antihypertensive medication. Average baseline values were systolic blood pressure 144.5 ± 24 mm Hg; diastolic blood pressure 79.3 ± 11 mm Hg; and mean blood pressure 101.0 ± 15 mm Hg. During the first month on D-OL-HDF, antihypertensive medication

was withdrawn and only one patient had high blood pressure. Reductions in blood pressure were observed although without achieving statistical significance (Table 4).

Cardiac parameters

Six patients had LVH assessments performed while on S-OL-HDF, just before the beginning of the study. A considerable reduction of LVH was observed when patients were switched to D-OL-HDF. LVM decreased from 167 ± 76 g at baseline to 124 ± 47 g after 3 months ($P < 0.01$) and 118 ± 37 g after 6 months ($P < 0.01$). LVMI decreased from 98 ± 45 g/m² at baseline to 72 ± 28 g/m² after 3 months ($P < 0.01$) and 69 ± 22 g/m² after 6 months ($P < 0.01$). LVH was present in only one patient at the end of the 6-month study period. A significant reduction in IVS, AWT, PWT, LWT, and IWT was also observed. There were no changes in LVDD, LVISD, LA, AO, LVEDV, LVESV, and EF (Table 6).

Finally, predialysis homocysteine levels decreased from 21.8 ± 3 μ mol/L on S-OL-HDF to 12.5 ± 2 μ mol/L after 3 months ($P < 0.01$) and 14.1 ± 5 μ mol/L after 6 months ($P < 0.01$) on D-OL-HDF.

DISCUSSION

Several groups [1–8] have reported excellent clinical results with higher frequency or daily dialysis because it is more physiologic and decreases the fluctuations in volume, solutes, and electrolytes. Improvement in comfort during and between dialysis sessions, clinical and biochemical parameters, anemia correction, hypertension control, nutrition status, and quality of life have been reported. OL-HDF is a safe technique that allows a considerable increase in convection, a high dialysis dose for both small and large molecules, and good control of anemia, nutrition, and blood pressure. In a previous paper [20], we reported that OL-HDF is the best technique for large molecule removal. The present study is the first experience reported in literature with D-OL-HDF and shows that the switch from standard three times a week

Table 4. Change from three times a week on-line hemodiafiltration (OL-HDF) to short daily on-line hemodiafiltration (D-OL-HDF). Hematologic parameters and blood pressure findings

	Baseline	Month 1	Month 2	Month 3	Month 4	Month 5	Month 6
Hemoglobin g/dL	12.3 ± 1	12.7 ± 1	12.5 ± 1	12.8 ± 1	12.6 ± 1	12.8 ± 1	12.8 ± 1
Hematocrit %	36.8 ± 5	37.8 ± 5	36.9 ± 5	38.2 ± 4	37.2 ± 3	38.0 ± 4	38.1 ± 4
Leucocytes ×10 ³ /μL	6.91 ± 1	6.91 ± 1	6.93 ± 2	6.33 ± 1	6.83 ± 1	6.85 ± 2	6.14 ± 1
Lymphocyte ×10 ³ /μL	1.93 ± 0.3	1.78 ± 0.5	1.71 ± 0.3	1.92 ± 0.4	1.87 ± 0.4	1.88 ± 0.3	1.84 ± 0.4
TS %	31.9 ± 12	31.2 ± 12	24.5 ± 7	31.7 ± 11	28.3 ± 10	28.4 ± 14	27.5 ± 11
Ferritin ng/mL	473 ± 263	360 ± 152 ^a	375 ± 160 ^a	312 ± 147 ^b	277 ± 151 ^b	273 ± 165 ^b	290 ± 177 ^b
Iron doses mg/week	24 ± 8	27 ± 14	28 ± 18	28 ± 18	37 ± 32	43 ± 33 ^a	46 ± 31 ^b
EPO doses IU/kg/week	48.0 ± 28	47.5 ± 28	47.4 ± 28	47.3 ± 28	47.0 ± 28	46.8 ± 28	46.8 ± 28
ERI	4.01 ± 2	3.83 ± 2	3.71 ± 2	3.68 ± 2 ^a	3.75 ± 2	3.76 ± 2	3.75 ± 2
SBP mm Hg	144 ± 24	140 ± 23	130 ± 29	131 ± 25	130 ± 17	131 ± 22	135 ± 18
DBP mm Hg	79.3 ± 11	78.3 ± 13	71.4 ± 14	70.1 ± 16	72.4 ± 11	75.2 ± 10	75.5 ± 12
MBP mm Hg	101 ± 15	98.9 ± 16	90.8 ± 18	90.4 ± 18	91.4 ± 12	93.8 ± 13	95.6 ± 14

Abbreviations are: TS, transferrin saturation; EPO, erythropoietin; ERI, EPO resistivity index (EPO doses/hemoglobin); SBP, systolic blood pressure; DBP, diastolic blood pressure; MBP, mean blood pressure.

^aP < 0.05; ^bP < 0.01 with respect to baseline value (ANOVA repeated measures); ^ciron dose was endovenous sodium ferric gluconate complex

Table 5. Change from three times a week on-line hemodiafiltration (OL-HDF) to short daily on-line hemodiafiltration (D-OL-HDF). Nutrition and inflammation findings

	Baseline	Month 1	Month 2	Month 3	Month 4	Month 5	Month 6
nPCR g/kg/day	0.931 ± 0.2	1.123 ± 0.3	1.075 ± 0.3	1.178 ± 0.3 ^a	1.160 ± 0.1 ^a	1.064 ± 0.3	1.129 ± 0.2
Body weight kg	67.8 ± 8	68.1 ± 8	68.1 ± 8	68.4 ± 8	68.8 ± 8	69.2 ± 8	69.4 ± 8
Weight gain kg	2.17 ± 0.5	1.28 ± 0.7 ^b	1.13 ± 0.4 ^b	1.24 ± 0.7 ^b	1.26 ± 0.4 ^b	1.26 ± 0.6 ^b	1.15 ± 0.6 ^b
Total protein mg/dL	7.0 ± 0.3	7.3 ± 0.6	6.8 ± 0.5	6.9 ± 0.3	7.0 ± 0.4	6.9 ± 0.4	7.0 ± 0.5
Albumin mg/dL	3.73 ± 0.2	3.74 ± 0.2	3.62 ± 0.3	3.77 ± 0.4	3.76 ± 0.2	3.65 ± 0.2	3.77 ± 0.2
Prealbumin mg/dL	33.1 ± 5	35.0 ± 6	32.3 ± 7	32.2 ± 7	33.8 ± 5	30.2 ± 5	32.3 ± 4
Transferrin mg/dL	169 ± 17	176 ± 26	162 ± 22	179 ± 15	184 ± 21	186 ± 20	195 ± 17
T cholesterol mg/dL	186 ± 44	200 ± 53	190 ± 59	203 ± 60	205 ± 66	203 ± 72	201 ± 58
HDL-c mg/dL	49.5 ± 8	57.0 ± 9	51.1 ± 16	58.9 ± 6	54.9 ± 7	56.3 ± 6	57.1 ± 11
LDL-c mg/dL	102 ± 33	113 ± 42	108 ± 38	115 ± 52	118 ± 56	135 ± 82	125 ± 45
Triglycerides mg/dL	165 ± 64	149 ± 96	152 ± 69	146 ± 82	157 ± 92	154 ± 62	162 ± 92
Fibrinogen mg/dL	428 ± 136	445 ± 35	486 ± 77	392 ± 24	420 ± 64	343 ± 38	353 ± 52
CRP mg/L	6.7 ± 7	8.4 ± 12	20.9 ± 36	3.8 ± 4	4.9 ± 7	5.4 ± 10	3.4 ± 4

Abbreviations are: nPCR, normalized protein catabolic rate; CRP, C-reactive protein; HDL-C, high-density lipoprotein cholesterol; LDL-C, low-density lipoprotein cholesterol.

^aP < 0.05; ^bP < 0.01 with respect to baseline value (ANOVA repeated measures)

Table 6. Change from three times a week on-line hemodiafiltration (OL-HDF) to short daily on-line hemodiafiltration (D-OL-HDF). Cardiac magnetic resonance imaging data

	Baseline	Month 3	Month 6
LVIDD mm	45.6 ± 2	47.1 ± 3	46.9 ± 4
LVISD mm	29.0 ± 3	29.2 ± 4	28.7 ± 3
IVS mm	15.0 ± 3.9	12.9 ± 3.2 ^a	12.7 ± 1.8 ^a
AWT mm	11.9 ± 4.6	8.3 ± 1.7 ^a	9.4 ± 1.4 ^a
PWT mm	11.9 ± 2.2	9.9 ± 2.6 ^b	9.8 ± 2.3 ^b
LWT mm	11.5 ± 2.9	9.7 ± 1.7	9.5 ± 2.0
IWT mm	12.6 ± 3.1	9.6 ± 2.7 ^b	10.1 ± 2.1 ^b
LA mm	34.7 ± 4.2	36.5 ± 5.0	37.6 ± 6.0
AO mm	33.8 ± 4.0	33.8 ± 2.6	33.8 ± 2.8
LVEDV mL	127.9 ± 18	116.4 ± 18	113.4 ± 13
LVESV mL	53.3 ± 9	39.7 ± 12	44.7 ± 11
EF %	60.7 ± 5	66.4 ± 10	62.3 ± 8
LVM g	166.9 ± 76	123.7 ± 47 ^b	118.1 ± 37 ^b
LVMi g/m ²	97.9 ± 45	72.4 ± 28 ^b	68.9 ± 22 ^b

Abbreviations are: LVIDD, left ventricular internal diameter at end-diastole; LVISD, left ventricular internal end-systole diameter; IVS, interventricular septal thickness; AWT, anterior wall thickness; PWT, posterior wall thickness; LWT, lateral wall thickness; IWT, inferior wall thickness; LA, left atrium dimension; AO, aortic root dimension; LVEDV, left ventricular end diastole volume; LVESV, left ventricular end systole volume; EF, ejection fraction; LVM, left ventricular wall mass; LVMi, LVM index.

^aP < 0.05; ^bP < 0.01 with respect to baseline value (ANOVA repeated measures)

OL-HDF to short D-OL-HDF has been satisfactory with excellent clinical tolerance and patient acceptance. It should be pointed out that, during the baseline period, the patients received a high dialysis dose and maintained a good control of anemia, nutrition, and blood pressure.

Although dialysis time was similar during both treatment schedules, an increase in the dialysis dose was obtained with D-OL-HDF, which confirms the beneficial effect of higher frequency. Although weekly spKt/V or weekly eKt/V were similar between both study periods, the EKR and standard Kt/V, proposed to measure the dialysis dose in dialysis regimens with different frequency, were 26% and 48% higher on D-OL-HDF, respectively. Weekly URR also was 52% higher in the daily schedule; this is a simple method that is able to show differences between regimens with different frequencies and the weekly reduction ratio can be used for any solute as well as urea.

Solute and fluid removal are the major goals of dialysis. Reduction of elevated predialysis uremic toxins levels may prevent or postpone the onset of dialysis-related complica-

tions. The difficulty in reducing large molecule plasma levels can be explained by the low distribution volume (approximately 20% of the body weight) or by a multi-compartmental model [21, 22]. In β_2 -microglobulin, the best results have been reported by Raj et al [23] with daily 8-hour nocturnal dialysis. They noted that the predialysis β_2 -microglobulin levels progressively declined from 27.2 to 13.7 mg/dL by 9 months. In the present paper, we observed a 21% reduction in predialysis plasma β_2 -microglobulin levels over a 6-month period.

Resistance to diffusion within tissues and organs creates solute disequilibrium gradients. The resistance to diffusion can be quantitated as the intercompartment mass-transfer coefficient (K_c) and is molecular size-sensitive. It is known that to obtain the same TAC in patients treated intermittently, the dialysis dose must be greater than during daily or continuous treatment. This phenomenon is magnified in solutes that have a lower K_c than urea [24, 25]. Our study supports this argument since we observed an increase in solute removal with higher frequency and this effect was more significant in solutes with greater molecular size or lower K_c (Fig. 3).

A decrease in erythropoietin dose has been reported by Vos, Zilch, and Kooistra [26], Galland et al [27], and Pierratos [28], amounting to 20%, 63%, and 40%, respectively. Buoncrisiani et al [1] initially reported an increase in hemoglobin from 8.1 ± 2 g/dL to 13.8 ± 3.8 g/dL, although the erythropoietin dose was not mentioned. We did not observe any changes in hemoglobin levels or erythropoietin dose in this study. Compared with these papers our patients received a higher dialysis dose with OL-HDF and baseline hemoglobin levels were higher with a low erythropoietin dose (<50 UI/kg/week). Ferritin was significantly reduced during the study and iron supplements were raised gradually to improve functional iron deficiency. The use of intravenous iron in daily dialysis experiences has not been clearly specified in the literature but it is possible that iron needs in daily dialysis may be higher. Kooistra et al [7] and Pierratos et al [5] did not observe any changes in erythropoietin dose and anemia control in initial reports but, in later papers, reported a decrease in erythropoietin dose related to changes in iron use [26, 28].

An increase in body weight (1.5 kg after 6 months) has been observed in this study as a marker of improvement in nutritional status. A rapid improvement in appetite was reported and an increase of nPCR was observed. Improvement of certain causes of anorexia such as post-dialysis fatigue, reduction in fluid overload, uremic milieu, medium and large-sized molecule removal was observed with the change to D-OL-HDF. Accumulation of substances <5000 D isolated from uremic plasma [29] and leptin [30, 31], 16,000 D have been related with anorexia and nutrition. Improvement in insulin resistance, growth hormone and insulin-like growth factor-1 (IGF-1) resis-

tance, and metabolic acidosis could be also involved [32]. Galland et al [27] reported similar data and found that short daily hemodialysis rapidly improves nutritional status, reporting an increase in body weight of 1.8 kg after 6 months without any changes in weekly Kt/V.

An improvement in phosphate control has been observed in this study and it has also been reported by most authors in daily dialysis treatment. Although we did not observe significant differences from predialysis phosphate levels, this could be due to the existence of confounding factors such as a reduction in phosphate binders (60% in our patients) and an increase in phosphate intake.

Cardiovascular disease is the most common cause of mortality in chronic hemodialysis patients, being the attributed cause of death in approximately 50% of the cases. In this study, the change to D-OL-HDF improved several cardiovascular risk factors such as arterial hypertension, hyperuricemia, LVH, and hyperhomocysteinemia. Hypertension is strongly associated with mortality and is very prevalent in dialysis patients ranging from 75% to 100%. Although our patients were relatively well controlled at baseline (only three patients were hypertensive and two were receiving drugs), better blood pressure control was achieved with D-OL-HDF without anti-hypertensive medications.

LVH is an independent cardiovascular risk factor and is strongly associated with mortality in dialysis patients [33]. LVH is present in 70% to 80% of the dialysis population [34]. Hypertension, poor extracellular fluid volume control, anemia, and uremia (accumulation of uremic toxins and circulatory vasoconstrictors) have been implicated in the pathophysiology of LVH. In the present study, we observed a marked regression of LVH. The LVMI was reduced by 30% after 6 months, after patients on S-OL-HDF were switched to D-OL-HDF. MRI has been validated for measurements of LV and RV wall mass [35]. Because of its high accuracy and superior reproducibility, MRI may be considered the gold standard for the assessment of ventricular dimensions, functional parameters and left and right ventricular mass [36]. Other authors have reported regression of cardiac hypertrophy measured with echocardiography. Fagugli et al [37] and Galland et al [27] reported a LVMI reduction of 18% and 31%, respectively, in patients receiving short daily dialysis. Recently, Chan et al [38] reported a LVMI reduction of 22% in 28 patients receiving slow nocturnal daily dialysis.

Hyperhomocysteinemia has been identified as an independent risk factor for cardiovascular disease [39]. The moderate elevation of total homocysteine concentrations in plasma, which is observed in at least 85% of hemodialysis patients, can be lowered by appropriate cofactor supplementation (folic acid, vitamin B₆, and B₁₂). However, administration of folic acid at pharmacologic doses,

either alone or combined with vitamin B₆ and B₁₂, is only partially effective in reducing plasma homocysteine and few dialysis patients normalize their homocysteine concentrations entirely [40]. This elevation of homocysteine concentrations is mainly due to the reduction of plasma homocysteine clearance, although the cause of this decrease is still unknown [41]. It is possible that the reduction of clearance is due to multiple abnormalities of the remethylation pathway that are not related to folate [42]. Therefore, the inhibition by uremic toxins of enzymes involved in the extrarenal homocysteine removal has been suggested as an alternative explanation for the elevated homocysteine levels in end-stage renal disease (ESRD) [43]. The effect of dialysis itself on plasma homocysteine concentration has been investigated in only a small number of studies. Interestingly, the reduction in homocysteine was not markedly different between high- and low-flux devices [44]. In the present study, plasma homocysteine levels decreased near normality without any changes in medical therapy, but only a higher frequency of dialysis. Similar data have been reported by Floridi et al [45], where homocysteine levels in 16 patients switched from standard hemodialysis to daily short hemodialysis diminished from 27.5 ± 11 $\mu\text{mol/L}$ to 17.2 ± 7 $\mu\text{mol/L}$.

CONCLUSION

The change from three times a week S-OL-HDF to short D-OL-HDF has been satisfactory. The main benefits observed in this study were the excellent clinical tolerance and patient acceptance, disappearance of post-dialysis fatigue, better dialysis adequacy (EKR, Kt/Vstd, weekly URR and TAD), a higher removal of medium and large-sized molecules, a reduction of phosphate binders, an improvement in nutritional status and, finally, a marked reduction of cardiovascular risk factors: better control of blood pressure without antihypertensive medication, reduction of ferritin and uric acid levels, regression of LVH and normalization of homocysteinemia levels.

ACKNOWLEDGMENTS

Part of this study was supported by grants from Spanish "Fondo de Investigaciones Sanitarias" (FIS 02/0811) and Fresenius Medical Care. Preliminary results from this study were presented at the 2002 National Congress of the Spanish Society of Nephrology, Bilbao, Spain.

Reprint requests to Francisco Maduell Canals, M.D., Servicio de Nefrología, Hospital General de Castellón, Av. Benicassim s/n., 12004 Castellón, Spain.
E-mail: maduell_fra@gva.es

REFERENCES

1. BUONCRISTIANI U, QUINTALIANI G, COZZARI M, et al: Daily dialysis: Long-term clinical metabolic results. *Kidney Int* 33(Suppl 24):S137–S140, 1988
2. HOMBROUCKX R, BOGAERT AM, LEROY F, et al: Limitations for short dialysis are the indications for ultra short daily auto dialysis. *Trans ASAIO* 35:503–505, 1989
3. MASTRANGELO F, ALFONSO L, NAPOLI M, et al: Dialysis with increased frequency of sessions (Lecce dialysis). *Nephrol Dial Transplant* 13(Suppl 6):139–147, 1998
4. PINCIAROLI AR: Results of daily hemodialysis in Catanzaro: 12 years experience with 22 patients treated for more than one year. *Home Int* 2:12–17, 1998
5. PIERRATOS A, OUWENDYK M, FRANCOEUR R, et al: Nocturnal hemodialysis: Three-year experience. *J Am Soc Nephrol* 9:859–868, 1998
6. DEPALMA JR, PECKER EA, MAXWELL MH: A new automatic coil dialyzer system for "daily" dialysis. *Proc Eur Dial Transplant Assoc* 6:26–34, 1969
7. KOOISTRA MP, VOS J, KOOMANS HA, VOS P: Daily home haemodialysis in The Netherlands: Effects on metabolic control, haemodynamics, and quality of life. *Nephrol Dial Transpl* 13:2853–2860, 1999
8. LOCKRIDGE RS, ANDERSON HK, COFFEY LT, CRAFT VW: Nightly home hemodialysis in Lynchburg, Virginia: Economic and logistic considerations. *Semin Dial* 12:440–447, 1999
9. CANAUD B, FLAVIER JL, ARGILES A, et al: Hemodiafiltration with on-line production of substitution fluid: Long-term safety and quantitative assessment of efficacy. *Contrib Nephrol* 108:12–22, 1994
10. MADUELL F, DEL POZO C, GARCIA H, et al: Change from conventional haemodiafiltration to on-line haemodiafiltration. *Nephrol Dial Transpl* 14:1202–1207, 1999
11. LORNOY W, BECAUS L, BILLIOUW JM, et al: On-line haemodiafiltration. Remarkable removal of β_2 -microglobulin. Long term observations. *Nephrol Dial Transpl* 15(Suppl 24):49–54, 2000
12. DAUGIRDAS JT: Second generation logarithmic estimates of single-pool variable volume Kt/V: An analysis of error. *J Am Soc Nephrol* 4:1205–1213, 1993
13. MADUELL F, GARCIA-VALDECASAS J, GARCIA H, et al: Validation of different methods to calculate Kt/V considering postdialysis rebound. *Nephrol Dial Transplant* 12:1928–1933, 1997
14. CASINO FG, LOPEZ T: The equivalent renal urea clearance: A new parameter to assess dialysis dose. *Nephrol Dial Transplant* 11:1574–1581, 1996
15. GOTCH FA: The current place of urea kinetic modelling with respect to different dialysis modalities. *Nephrol Dial Transplant* 13(Suppl 6):10–14, 1998
16. LOPOT F, VÁLEK A: Time-averaged concentration—Time-averaged deviation: A new concept in mathematical assessment of dialysis adequacy. *Nephrol Dial Transplant* 3:846–848, 1998
17. BERGSTRÖM J, WEHLE B: No change in corrected β_2 -microglobulin concentration after cuprophane haemodialysis. *Lancet* 1:628–629, 1987
18. SKLAR A, NEWMAN N, SCOTT R, et al: Identification of factors responsible for postdialysis fatigue. *Am J Kidney Dis* 34:464–470, 1999
19. RADEMAKERS FE, BOGAERT J: Cardiac function, in *Magnetic Resonance of the Heart and Great Vessels*, edited by BOGAERT J, DUERINCKX AJ, RADEMAKERS FE, Berlin, Springer, 1999, pp 95–111
20. MADUELL F, NAVARRO V, CRUZ MC, et al: Osteocalcin and myoglobin removal in on-line hemodiafiltration versus low and high-flux hemodialysis. *Am J Kidney Dis* 40:582–589, 2002
21. ODELL RA, SŁOWIACZEK P, MORAN JE, SCHINDHELM K: Beta₂-microglobulin kinetics in end-stage renal failure. *Kidney Int* 39:909–919, 1991
22. VINCENT C, REVILLARD JP, GALLAND M, TRAEGER J: Serum beta₂-microglobulin in hemodialysed patients. *Nephron* 21:260–269, 1978
23. RAJ DSC, OUWENDYK M, FRANCOEUR R, PIERRATOS A: β_2 -Microglobulin kinetics in nocturnal haemodialysis. *Nephrol Dial Transplant* 15:58–64, 2000
24. DEPNER TA: Quantifying hemodialysis and peritoneal dialysis: Examination of the peak concentration hypothesis. *Semin Dial* 7:315–317, 1994
25. DEPNER TA: Benefits of more frequent dialysis: Lower TAC at the same Kt/V. *Nephrol Dial Transplant* 13(Suppl 6):20–24, 1998
26. VOS PF, ZILCH O, KOOISTRA MP: Clinical outcome of daily dialysis. *Am J Kidney Dis* 37(Suppl 2):S99–S102, 2001
27. GALLAND R, TRAEGER J, ARROUCHE W, et al: Short daily hemodialy-

- sis and nutritional status. *Am J Kidney Dis* 37(Suppl 2):S95–S98, 2001
28. PIERRATOS A: Nocturnal home haemodialysis: An update on a 5-year experience. *Nephrol Dial Transplant* 14:2835–2840, 1999
29. ANDERSTAM B, MAMOUN A, SÖDERSTEN P, BERSTRÖM J: Middle-sized molecule fractions isolated from uremic ultrafiltrate and normal urine inhibit ingestive behavior in the rat. *J Am Soc Nephrol* 7:2453–2460, 1996
30. YOUNG GA, WOODROW G, KENDALL S, et al: Increased plasma leptin/fat ratio in patients with chronic renal failure: A cause of malnutrition? *Nephrol Dial Transplant* 12:2318–2323, 1997
31. MERABET E, DAGOGO-JACK S, COYNE DW, et al: Increased plasma leptin concentrations in end stage renal disease. *J Clin Endocrinol Metab* 82:847–850, 1997
32. IKIZLER TA, HIMMELFARB J: Nutritional complications in chronic hemodialysis and peritoneal dialysis patients, in *Complications of Dialysis*, edited by LAMEIRE N, METHAS RL, New York, Marcel Dekker, Inc., 2000, pp 405–425
33. SILBERBERG JS, BARRE PE, PRICHARD SS, SNIDERMAN AD: Impact of left ventricular hypertrophy on survival in end-stage renal disease. *Kidney Int* 36:286–290, 1989
34. FOLEY RN, PARFREY PS, HARNETT JD, et al: Clinical and echocardiographic disease in patients starting end-stage renal disease therapy. *Kidney Int* 47:186–192, 1995
35. PATTYNAMA PM, LAMB HJ, VAN DER VELDE EA, et al: Left ventricular measurements with cine and spin-echo MR imaging: A study of reproducibility with variance component analysis. *Radiology* 187:261–268, 1993
36. HIGGINGS CB: Which standard has the gold? *J Am Coll Cardiol* 19:1608–1609, 1992
37. FAGUGLI RM, REBOLDI G, QUINTALIANI G, et al: Short daily hemodialysis: Blood pressure control and left ventricular mass reduction in hypertensive hemodialysis patients. *Am J Kidney Dis* 38:371–376, 2001
38. CHAN CT, FLORAS JS, MILLER JA, et al: Regression of left ventricular hypertrophy after conversion to nocturnal hemodialysis. *Kidney Int* 61:2235–2239, 2002
39. BOSTOM AG, SHEMIN D, LAPANE KL, et al: Hyperhomocysteinemia and traditional cardiovascular risk factors in end-stage renal disease patients on dialysis: A case control study. *Atherosclerosis* 114:93–103, 1995
40. MASSY ZA: Importance of homocysteine, lipoprotein (a) and non-classical cardiovascular risk factors (fibrinogen and advanced glycation end-products) for atherogenesis in uraemic patients. *Nephrol Dial Transplant* 15(Suppl 5):81–91, 2000
41. GUTTORMSEN AB, ELAND PM, SVARSTAD E, REFSUM H: Kinetic basis of hyperhomocysteinaemia in patients with chronic renal failure. *Kidney Int* 52:495–502, 1997
42. VAN GULDENER C, STAM F, STEHOUEWER CD: Homocysteine metabolism in renal failure. *Kidney Int* 59(Suppl 78):S234–S237, 2001
42. ARNADOTTIR M, BERG AL, HEGBRAND J, et al: Influence of hemodialysis on plasma total homocysteine concentration. *Nephrol Dial Transplant* 14:142–146, 1999
44. BIASIOLI S, SCHIAVON R, PETRONISO L, et al: Dialysis kinetics of homocysteine and reactive oxygen species. *ASAIO J* 44:423–432, 1998
45. FLORIDI A, BUONCRISTIANI U, FAGUGLI RM, et al: Daily hemodialysis effectively lowers hyperhomocysteinemia in uremic patients. *J Am Soc Nephrol* 9:233A, 1998